

ACTIVE INGREDIENTS

Parathion (O,O-Diethyl O-p-nitrophenyl thiophosphate).....	2.0%
INERT INGREDIENTS.....	93.0%
Total.....	100.0%



POISON

ANTIDOTE

WARNING: Keep Out of Reach of Children

Poisonous if swallowed or inhaled or absorbed through skin or eyes. Rapidly absorbed through skin. Do not get in eyes or on skin or clothing. In case of contact, wash immediately with soap and water. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type approved by the U.S. Bureau of Mines for Parathion protection. Wash arms, hands and face thoroughly after handling and before eating or smoking. In case of contact wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before reuse. Keep away from animals, feed and foodstuffs. Do not feed to dairy animals or to animals being furnished for slaughter. Keep unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.

Not for use or storage near around the home.
 Highly toxic to fish and wildlife. Birds and other wildlife in treated areas may be harmed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.
 Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS

As recommendations for the use of Parathion products vary in different agricultural areas, it is suggested that you consult your State Agricultural Experiment Station before using this product. If the crop to be dusted is for canning purposes consult your cannery fieldman before dusting. General recommendations for the use of this dust are listed below and are believed to be of value but in no way guaranteed.

BEANS - To control Aphids, Red Spiders and Mexican Bean Beetles apply dust at the rate of 15 to 20 lbs. per acre when insects appear and repeat as necessary. Do not use dust within 15 days of harvest.

CABBAGE, CAULIFLOWER and BROCCOLI - To control Imported Cabbage Worms and Aphids apply dust at the rate of 20 lbs. per acre when insects appear and repeat at 7 to 10 day intervals as needed. Do not apply dust on cauliflower and broccoli within 7 days of harvest. Do not treat cabbage within 10 days of harvest.

CARROTS - To control Six Spotted Leafhopper apply dust at the rate of 15 to 20 lbs. per acre and repeat at 7 day intervals if necessary. To be effective the field margins should be treated also. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed.

CELERY and LETTUCE - To control Aphids, Six Spotted Leafhopper and Tarnish Plant Bugs apply at the rate of 20 lbs. per acre. For leafhoppers be sure to treat margins also. Do not use dust within 21 days of harvest for celery and lettuce.

ONIONS (dry onions only) To control Onion Maggot apply 25 lbs. per acre to the open furrow immediately following seeding but before the furrow is closed. 14 days later apply dust at the rate of 15 to 20 lbs. per acre and repeat every 3 to 5 days as necessary. Do not use within 15 days of harvest.

PEAS - To control Pea Aphids apply dust at the rate of 15 to 20 lbs. per acre when insects first appear. Do not apply within 10 days of harvest.

POTATOES - To control Grasshoppers apply dust at the rate of 15 to 20 lbs. per acre when they increase in numbers and are not affected by regular dust or spray schedule. Do not use on potatoes within 5 days of harvest.

NON-WARRANTY

Sella makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595 249

9-21-71
 595-249

Net Wt. 50 Lbs.