

(CONTAINS 5% SEVIN*)

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS:	<u>95.0%</u>
Total	100.0%

*Sevin is the registered Trade Mark of Union Carbide Corp. for the active ingredient 1-naphthyl N-methylcarbamate.

DIRECTIONS

As conditions may vary in local agricultural areas it is recommended that your State Agricultural Experiment Station be consulted for additional recommendations in each area.

BLUEBERRIES – To control Fruitworms and Fruit Fly apply Haviland 5-S Dust at the rate of 40 lbs. per acre. Make first application immediately after bloom or as soon as insects become active and repeat at 10-day intervals until fruit is harvested.

STRAWBERRIES – To control Spittlebug and Leafrollers apply this dust at the rate of 40 lbs. per acre for full coverage in regular cover applications. Do not apply within 1 day of harvest.

GRAPES – To control Grape Leaf Folder, Leafhopper and Berry Moth apply at the rate of 40-60 lbs. per acre. Make first application just before first brood leaf folder larvae emerge and repeat at weekly intervals as necessary.

TOMATOES, EGGPLANT and PEPPERS – To control European Corn Borer, Fall Armyworm, Lace Bug, Tomato Fruitworm, Tomato Hornworm, Stinkbugs, Tarnished Plant Bug apply 25-40 lbs. of dust per acre. Make first application when insects or their damage first appear and repeat at 7-10 day intervals as needed.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS and KOHLRABI – To control Armyworms, Flea Beetle, Harlequin Bug and Corn Earworm apply this dust at the rate of 35-40 lbs. per acre when insects or their damage appear and repeat at 5 to 7 day intervals as necessary. Do not make application within 3 days of harvest.

CAUTION:

Keep Out of Reach of Children

Harmful if swallowed. Avoid contact with skin. Wash thoroughly after using. Avoid breathing dust. Avoid contamination of feed and foodstuffs. Use only on recommended crops and at recommended rates. Avoid storage near feed or foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

For protection of bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn bee keeper well in advance to locate hives at a safe distance until 1 week after application.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-252

Net Wt. 50 Lbs.