

PARATHION MANEB DUST NO. 27

ACTIVE INGREDIENTS:

Parathion (O,O-Diethyl O-p-Nitrophenyl thiophosphate) 2.0%

Maneb (Manganese ethylene bisdithiocarbamate) 7.0%

INERT INGREDIENTS: 91.0%

100.0%

Recommended for control of various insects and fungus diseases on Vegetable Crops as listed below.

DIRECTIONS

As recommendations for the use of Parathion products vary in different agricultural areas, it is suggested that you consult your State Agricultural Experiment Station before using this product. If the crop to be dusted is for canning purposes, consult your cannery fieldman before dusting. This dust should be applied at the rates suggested below depending on the size and spacing of the plants and the method of application.

CARROTS — To control Aphids, Six Spotted Leafhopper, Alternaria Leaf Blight and Cercospora Blight apply 20 to 30 lbs. per acre. Begin application when plants are coming up and repeat weekly as long as insects or blight threatens. In dusting for leafhopper treat both field and field-margins. Do not use on carrots within 15 days of harvest (21 days if treated tops are to be used for feed or food).

CELERY — To control Aphids, Six Spotted Leafhopper, Tarnish Plant Bug, Early and Late Blight use 30-40 lbs. per acre weekly. Begin dusting as soon as insects appear or blight threatens. Do not use after celery begins to bunch or is half grown. Remove excess residues by stripping, trimming, and washing.

CUCUMBER, MELON and SQUASH — To control Cucumber Beetle, Aphids, Stink Bug, Melonworm, Pickleworm, Anthracnose, Cercospora leaf spot and Downy Mildew apply 20 to 25 lbs. per acre when insect appears or disease threatens. Do not use on cucumbers before vining. Do not use within 15 days of harvest on cucumbers or squash nor within 7 days of harvest on melons.

ONIONS — To control Downy Mildew, Blast (Botrytis leaf blight) and Purple Blotch apply 30 to 40 lbs. per acre. For fungus control use at weekly intervals when disease is reported in the area. Do not apply within 15 days of harvest.

POTATOES — To control grasshoppers and Early and Late Blight apply 30 to 50 lbs. per acre when insects appear or when blight threatens. Do not apply within 5 days of harvest.

NOTE: For disease control closer to harvest than the given limitations a suitable fungicide should be used alone.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DANGER: Keep Out of Reach of Children
MAY BE FATAL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN! Do not get on skin, in eyes, or on clothing. Wear

natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Parathion protection. Keep all unprotected persons out of operational areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Not for use or storage in or around the home.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Toxic to fish and wildlife. Do not contaminate any body of water nor apply when weather conditions favor drift from the area treated. Do not apply to any area not specified on the label. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.



POISON



Antidote — Internal: Give a teaspoon of mustard in a glass of water. Repeat until gastric fluid is clear. Have victim lie down and remain quiet after the physical vomiting has been carried out. Call a physician immediately. Atropine Sulfate (in advice of physician) 2 PAM is also antidotal and may be administered in conjunction with atropine.

External: If skin is contaminated, wash thoroughly with soap and water.

USDA Reg. No. 595-256

Net Wt. 50 Lbs.

8-18-71
595-256