

(INSECTICIDE-FUNGICIDE DUST)

ACTIVE INGREDIENTS:

Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate)	1.0%
Metallic Copper (From Tribasic Copper Sulfate)	3.5%
Maneb (Manganese ethylenebisdithiocarbamate)	3.5%
INERT INGREDIENTS:	92.0%
	100.0%



POISON



ANTIDOTE

Internal: Give a teaspoon of mustard in a glass of warm water. Repeat until vomit begins. Then lie on your side and remain quiet after the physical vomiting has been gained. Call a physician immediately. Atropine Sulfate on advice of physician. DIPAM is also antidotal and may be administered in conjunction with atropine.

External: If skin is contaminated, wash thoroughly with soap and water immediately.

DANGER: Keep Out of Reach of Children

Poisonous if swallowed, inhaled or absorbed through skin or eyes. Rapidly absorbed through skin! Do not get in eyes or on clothing. In case of contact, wash immediately with soap and water. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Parathion protection. Wash arms, hands and face thoroughly after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep away from children, animals, feed and foodstuffs. Do not feed to dairy animals or to animals being furnished for slaughter. Keep unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.

Highly toxic to fish and wildlife. Do not contaminate any body of water, nor apply when conditions favor drift from the area treated. Do not apply to any area not specified on the label. Birds feeding on treated areas may be killed.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Not for use or storage in or around the home.

DIRECTIONS

As recommendations may differ in various agricultural areas, it is suggested that you consult your local State Agricultural Extension Service or Agricultural Experiment Station for specific rates and timing of applications. If the crop to be dusted is for canning purposes, consult your cannery fieldman before dusting. This dust should be applied at the rates suggested below depending on the size and spacing of the plants and the method of application.

CARROTS: To control Aphids, Six Spotted Leafhopper, Alternaria Leaf Blight, and Cercospora Blight apply 30 to 40 lbs. per acre. Begin application when plants are coming up and repeat weekly as long as insects or blight threaten. Do not use on carrots within 15 days of harvest (21 days if treated tops are to be used for feed or food).

CELERY: To control Aphids, Six Spotted Leafhopper, Tarnished Plant Bug and Early and Late Blight — apply 30 to 40 lbs. per acre weekly. Begin dusting as soon as insects or disease threaten. Do not use after celery begins to bunch or is half grown. Remove excess residues by stripping, trimming and washing.

CUCUMBER, MELON AND SQUASH: To control Cucumber Beetle, Aphids, Stink Bug, Melonworm, Pickleworm, Anthrachnose, Cercospora Leaf Spot and Downy Mildew - apply 30 to 50 lbs. per acre when insects or disease threatens and repeat at weekly intervals as necessary. Do not use on cucumbers before vining or within 15 days of harvest. Do not use on melons within 7 days or squash within 15 days of harvest.

ONIONS: To control Downy Mildew, Blast (Botrylis Leaf Blight) and Purple Blotch — apply 30 to 40 lbs. per acre. For disease control apply at weekly intervals when disease threatens. Do not apply within 5 days of harvest.

POTATOES: To control Grasshoppers, Early and Late Blight, apply 30 to 50 lbs. per acre when insects appear or when blight threatens. Do not apply within 5 days of harvest.

NOTE: For disease control closer to harvest than the given limitations above, a suitable fungicide should be used alone.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.