

REST OF THE PRODUCT AVAILABLE

HAYILAND ATRAZINE LINURON WEED KILLER

**For Selective Weed Control In Corn
(Wettable Powder)**

REPT
1969
595-268

- Active Ingredients:
- Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine) .. 30.8%
 - Linuron [3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea] 30.8%
 - Inert Ingredients: 38.4%

U.S. Patent No. 3,022,150
U.S.D.A. Reg. No. 595-268

CAUTION!

HARMFUL IF SWALLOWED. MAY IRRITATE EYES, NOSE, THROAT AND SKIN.
Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Avoid contamination of feed and foodstuffs.

Keep Out of Reach of Children

Net Wt. 5 Lbs.



BEST AVAILABLE

4

HAVILAND ATRAZINE LINURON WEED KILLER

IMPORTANT

Use HAVILAND ATRAZINE LINURON WEED KILLER only as specifically recommended. Over-dosage, caused by inaccurate preparation of spray mixture, improper calibration of equipment or application at a speed slower than used for calibration, should be avoided as injury to crop may result. Do NOT apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of dry powder or spray to desirable plants. Do NOT contaminate domestic waters. Keep from contact with fertilizers, insecticides, fungicides, and seed. Thoroughly clean all traces of HAVILAND ATRAZINE LINURON WEED KILLER from application equipment immediately after use; otherwise, crop injury may result when equipment is used again. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

DIRECTIONS

HAVILAND ATRAZINE LINURON WEED KILLER is a wettable powder to be mixed in water and applied as a spray for selective control of annual weeds and grasses in field corn. It is non-corrosive, non-volatile and non-flammable. Treatment provides control of germinating annual weed seedlings such as foxtail (including giant foxtail), barnyardgrass, crabgrass, lambsquarter, pigweed, purslane, mustard, ragweed and smartweed. Treatment will not control established quackgrass, nutgrass, Johnsongrass, bermudagrass, or other perennial weeds.

HOW TO USE

Make a single application as a band (over the row) or broadcast spray after planting but before crop emerges. For accuracy and convenience, planting and spraying should be combined in one operation. Apply with a properly calibrated fixed boom power sprayer, using sufficient water (equivalent to 25 to 40 gallons per acre on a broadcast basis) to provide thorough and uniform coverage of the ground. Avoid overlapping, and shut off spray boom while starting, turning, slowing or stopping, or injury to the crop may result. Continuous agitation is required to keep the material in suspension. Best results are obtained when HAVILAND ATRAZINE LINURON WEED KILLER is moved by moisture (rainfall or overhead irrigation) into the upper layer of soil within 2 weeks after application. Lack of sufficient moisture (1 to 2 inches is generally sufficient) following treatment will generally result in unsatisfactory weed control. Soils should be well prepared and free as possible from trash and clods. Seed should be planted to a depth of at least 1 1/2 inch or crop injury may result. If soil becomes crusted before corn emerges, a shallow rotary hoeing may be made. If rainfall is insufficient to activate the chemicals following treatment, rotary hoeing or standard cultivation should be used before weeds become well established; otherwise, avoid cultivating or disturbing the sprayed area, such as covering treated band with soil while cultivating middles. Fields treated with HAVILAND ATRAZINE LINURON WEED KILLER may be planted to the same crop if initial seeding fails to germinate or produce a stand. Thoroughly rework soil before planting. Do not retreat with a second application during the same crop year as injury to the crop may result. Do not replant treated areas with crops other than corn within 6 months after application or crop may be injured. Do not follow treated corn with sugar beets, tobacco or vegetables in rotation. Prior to replanting to any crop following treated corn, thorough seedbed preparation including fall or spring plowing is recommended.

APPLY THE SPRAY AT THE FOLLOWING RATES

Soil Type*	Lb. HAVILAND ATRAZINE LINURON WEED KILLER in 25 to 40 gallons water per acre (if ground actually sprayed).	Example — Band treatment lbs. crop area 14" Band on 42" Rows.
Light colored—Low organic matter 2% or less		
Sandy loam	1.6	5
Silt loam	2.0	7
Clay loam	2.5	8
Dark colored—moderate organic matter		
Silt loam	4.0	13
Clay loam	5.0	17

* Do not use on sand or leamy sand soils as crop injury may result.

EQUIPMENT: Use a tractor-mounted power sprayer properly calibrated to a constant speed and rate of delivery (see CALIBRATION section). Sprayer should be designed to treat the same number of rows as the planter used. Make sure spray equipment is clean (free of scale, rust, dirt, oil and pesticide deposits). Screens should be 50-mesh or coarser. Booms and hoses should be maximum of 3/4 inch diameter. MATERIALS MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION. Agitate by mechanical or hydraulic means. With hydraulic system, use a separate agitation line in addition to by-pass line; both lines should terminate at bottom of tank to minimize foaming and to give maximum agitation. Spray booms must be shut off while starting, turning, slowing or stopping, or injury to crop may result.

CALIBRATION: The following is designed for application of indicated number of gals. spray per acre of ground actually sprayed; it is based on width of band treated and is not affected by row spacing.

Step 1: To calibrate, use tractor gear and throttle setting which will be used for actual application (full throttle is preferred to insure maximum pump capacity). With only water in the tank, determine number of seconds required for tractor to travel in the field the measured distance shown below — opposite desired band width and under proper spray volume.

Band Width	GALS. SPRAY PER ACRE OF GROUND ACTUALLY SPRAYED				Band Width	GALS. SPRAY PER ACRE OF GROUND ACTUALLY SPRAYED			
	25	30	35	40		25	30	35	40
10 in.	523 ft.	436 ft.	373 ft.	326 ft.	16 in.	327 ft.	272 ft.	233 ft.	204 ft.
12 in.	436 ft.	363 ft.	311 ft.	272 ft.	18 in.	291 ft.	242 ft.	207 ft.	181 ft.
14 in.	374 ft.	311 ft.	266 ft.	232 ft.	20 in.	262 ft.	218 ft.	187 ft.	163 ft.

Adjust height of boom to obtain desired band width. With only water in the tank and using the same throttle setting as in Step 1 but with tractor standing still, adjust pressure so that EACH NOZZLE delivers ONE QUART or spray in the exact number of seconds as determined in Step 1 for the selected band width.

Step 2: Equipment is now ready to apply the required spray volume, provided nozzle height, gear and throttle setting are not changed.

NOTICE TO BUYER: SELLER makes no warranty of any kind, express or implied, concerning the use of this product. BUYER assumes all risk of use or handling whether in accordance with directions or not.

HAVILAND AGRICULTURAL CHEMICAL COMPANY
GRAND RAPIDS, MICHIGAN

7