

BOTRAN 6% DUST
(Fungicide)

ACCEPTED

APR 23 1966

ACTIVE INGREDIENTS

*Botran (2,6-dichloro-4-nitroaniline)

6.6%

INERT INGREDIENTS

93.4%
100.0%

*Botran is a registered trademark of the TUCO Products Company.

Botran has been found compatible with formulations of Captan, Zineb, DDT, Guthion, Parathion, Thiodan, and Carbaryl.

DIRECTIONS

PEACHES---To control Blossom Blight apply at the rate of 60 lbs. per acre at pink bud and full bloom. To control Brown Rot and Rhizopus Rot apply at the rate of 60 lbs. per acre 18 days, 10 days, and 1 day prior to harvest. Do not apply less than 1 day prior to harvest.

STRAWBERRY---To control Botrytis Rot and Rhizopus Rot apply at the rate of 25 lbs. per acre. Begin dusting when new leaves are expanded and blossom buds visible; repeating applications weekly through harvest.

SWEET CHERRY---To control Blossom Blight apply at the rate of 50 lbs. per acre at popcorn bloom, full bloom, and petal fall. To control Brown Rot apply at the rate of 50 lbs. per acre at 10 days and 1 day prior to harvest.

GRAPES---To control Botrytis Rot apply at the rate of 30 lbs. per acre. Begin application while clusters are still loose and repeat at two-week intervals. Do not apply within 1 day of harvest.

TOMATOES (Greenhouse)---To control Botrytis (stem canker), apply at the rate of 30 lbs. per acre. Dust stem of plant from ground level up to a height of 18 to 24 inches. Repeat weekly as necessary.

LEAF LETTUCE (Greenhouse)---To control Botrytis Rot apply at the rate of 30 lbs. per acre. Apply dust 10 days following transplanting; repeating when plants are half mature. Do not apply to wilted plants, seedlings or within 10 days of harvest.

ROSE, GLADIOLUS, CHRYSANTHEMUM, GERANIUM---To control blight (Botrytis) apply at the rate of 30 lbs. per acre. In the field, apply prior to shipping and storage. When storing, thoroughly dust storage area and all canes and dead foliage; repeat application during long storage periods. For stock cuttings or greenhouse plants, begin dusting when disease is anticipated or first appears. Dust foliage and flowers at 10 to 14 day intervals. During prolonged damp, cold periods, apply every 7 to 10 days.

CLAS. To control Blight (Botrytis) apply at the rate of 30 lbs. per acre, dusting thoroughly. Force fungicide between spikes. Apply to plants and cut flower spikes after harvest and before shipment or storage.

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not breathe dust. Avoid contact with skin and eyes. In case of contact, remove contaminated clothing and wash skin with soap and water; for eyes, flush with water for 15 minutes and obtain medical attention. Do not contaminate feed or foodstuffs. Do not feed treated forage to livestock.

NO WARRANTY

The use and handling of this material being beyond our knowledge and control, we so make any warranty expressed or implied as to the effects of such use, whether or not in accordance with directions or claimed to be so.

Net Wt. 50 Lbs.

Net Wt. 50 Lbs.

HAVILAND AGRICULTURAL CHEMICAL COMPANY

GRAND RAPIDS, MICHIGAN