

# thiodan<sup>®</sup> zineb dust 36

## ACTIVE INGREDIENTS:

Endosulfan (Hexachlorohexahydromethano  
2, 4, 3 Benzodioxathiopin oxide) ..... 3.0%  
Zineb (Zinc Ethylene bisdithiocarbamate) ..... 6.0%

INERT INGREDIENTS: ..... 91.0%

Total ..... 100.0%

<sup>5</sup>Thiodan is a registered trademark of Farbwerke Hoechst A.G. for the active ingredient commonly known as Endosulfan.

## DIRECTIONS

The following directions are furnished by us and are believed to be reliable but in no way guaranteed. Make first application when insects appear or disease threatens and repeat weekly if necessary. Consult your State Agricultural Experiment Station or Extension Service for specific timing and dosage rates in your area.

**BEANS:** To control Mexican bean beetles, anthracnose, downy mildew, and rust, apply at the rate of 25 to 30 lbs. per acre. Do not apply after pods begin to form. Do not apply more than 3 times per season. Do not use on lima beans that are to be removed from the field for processing. Do not feed treated threshings to livestock.

**BROCCOLI, CABBAGE AND CAULIFLOWER:** To control cabbage looper, imported cabbage worm, Diamond-Back moth larvae, alternaria leaf blight and downy mildew, apply at the rate of 25 to 30 lbs. per acre. Do not apply on broccoli and cabbage within 7 days of harvest. Do not apply on cauliflower within 14 days of harvest.

**CUCUMBERS, MELONS AND SQUASH:** To control aphids, anthracnose, alternaria blight, and downy mildew, apply at the rate of 25 to 30 lbs. per acre. Apply as needed up to harvest.

**HEAD LETTUCE:** To control aphids, cabbage looper, imported cabbage worm, and Diamond-Back moth larvae, apply at the rate of 35 to 40 lbs. per acre. To control downy mildew, apply at the rate of 25 to 35 lbs. per acre. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not apply within 14 days of harvest. Do not feed crop refuse to livestock.

**PEPPERS AND TOMATOES:** To control aphids, Colorado potato beetles, flea beetles, anthracnose, early and late blight, apply at the rate of 15 to 20 lbs. per acre. Do not apply within 1 day of harvest.

## CAUTION: Keep Out of Reach of Children

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe dust or spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, obtain medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. Wear a mask or respirator of a type approved by the U.S. Bureau of Mines for Endosulfan protection.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when conditions favor drift from the area treated. Do not contaminate feed or foodstuffs.

Apply this product only as specified on this label.

Do not reuse empty container. Destroy when empty by burying with waste or burning. Stay away from smoke or fumes.

## NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 595-274

6-13-72 with comment,  
595-274

Net Wt. 50 Lbs.