

ACTIVE INGREDIENTS:

Captan [N- [(trichloromethyl)] thio-4-cyclohexene-1, 2-dicarboximide]	7.5%
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS	87.5%
	100.0%

*Sevin is the registered trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

DIRECTIONS

APPLES: To control apple scab, Brooks fruit spot, black pox, and botryosphaeria, apply at the rate of 25 to 35 lbs. per acre. To control codling moth, aphids, apple maggot, plum curculio, Eastern tent caterpillar, San Jose scale crawlers and Forbes scale crawlers, apply at the rate of 70 to 120 lbs. per acre. Begin with second cover and continue at 10-14 day intervals as necessary. Do not apply within 1 day of harvest.

CHERRIES: To control brown rot and leaf spot, apply at the rate of 25 to 50 lbs. per acre. To control codling moth, Forbes scale crawlers, San Jose scale crawlers, and red banded leaf rollers, apply at the rate of 80 to 120 lbs. per acre. Begin application at petal fall and repeat at 10 day to 14 day intervals through cover applications. Do not apply within 1 day of harvest.

PEACHES: To control brown rot, peach scab, plum curculio, and peach twig borer, apply at the rate of 30 to 60 lbs. per acre. To control codling moth and San Jose scale crawlers, apply at the rate of 80 lbs. per acre. Begin at petal fall and repeat at 10 to 14 day intervals through cover applications. Do not apply within 1 day of harvest.

CUCUMBERS: To control angular leaf spot, anthracnose, downy mildew, and pickleworm, apply at the rate of 15 to 20 lbs. per acre. To control cucumber beetles, squash bugs, flea beetles, and leaf hopper, apply at the rate of 20 to 25 lbs. per acre. Apply when insects or disease threaten and repeat as necessary.

TOMATOES: To control alternaria, late blight, septoria, stemphylium, tarnished plant bugs, tomato fruitworm, stink bugs, tomato hornworm, European corn borer, and leaf hoppers, apply at the rate of 30 to 40 lbs. per acre. Begin when insects or disease threatens and repeat at 7 to 10 day intervals as necessary.

POTATOES: To control Colorado potato beetles, flea beetles, leaf hoppers, early blight (alternaria), and late blight, apply at the rate of 30 to 40 lbs. per acre. Begin application when insects or disease threatens and repeat as necessary.

CAUTION: Keep Out of Reach of Children

Harmful if swallowed or inhaled. Avoid breathing dust. Wear a respirator or mask of a type approved by the U. S. Bureau of Mines. Avoid skin contact. Wash thoroughly after using and before eating or smoking. Avoid contamination of feed and foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative agricultural extension service.

This product is toxic to fish. Keep out of any body of water. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of waste. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. Reg. No. 595-282

Net Wt. 50 Lbs.



HAYILAND

1400 BERLING N.W. GRAND RAPIDS, MICHIGAN 49502

ACCEPTED WITH COMMENTS