diazinon* maneb dust no. 27

ACTIVE INGREDIENTS:

(O. O-diethyl O-(2 -isopropyl-4-methyl- 6-			
pyrimidinyl)] thiophosphate	 	 	. 2.0%
Maneb (Manganese ethylenebisdithiocarbamate)	 	 	.7.0%
INERT INGREDIENTS:	 	 9	91.0%
Tasal		1/	00.00/
Total	 	 !!	JU.U%

*Diazinon is a proprietary material of Geigy Agricultural Chemical Co. for the active material O, O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) thiophosphate.

DIRECTIONS

The following directions are furnished by us and are believed to be of value but in no way guaranteed. This dust should be applied at the rates of 30 to 50 lbs. per acre unless otherwise specified, depending on the size and spacing of the plants and the method of application. Make first application when insects appear or disease threatens and repeat at 7 to 10 day intervals, as necessary.

BEANS — To control Bean Aphids, Mexican Bean Beetles, Leafhoppers, Downy Mildew and Rust, apply at the rate of 30 to 45 lbs. per acre. Do not apply within 7 days of harvest. (After harvest is completed, treated forage may be fed to livestock after 1 day following treatment or cut for hay after 4 days following treatment.)

BROCCOLI, CABBAGE AND CAULIFLOWER — To control Aphids, Imported Cabbage worms, Downy Mildew and Alternaria Leaf Spot apply at the rate of 30 to 35 lbs. per acre. Do not apply to Cabbage within 7 days of harvest nor to Broccoli and Cauliflower within 5 days of harvest.

CUCUMBERS, MELONS AND SQUASH — To control Aphids, Leafhoppers, Downy Mildew, and Anthracnose. Do not apply to Summer Squash or Cucumbers within 7 days of harvest or to Winter Squash or Melons within 3 days of harvest.

ONIONS — To control Onion Thrips, Botrytis Leaf Blight, Downy Mildew and Purple Blotch apply at the rate of 30 to 40 lbs. per acre. Do not apply within 10 days of harvest.

POTATOES — To control Aphids, Colorado Potato Beetles, Flea Beetles, and Early and Late Blight. Do not apply within 35 days of harvest.

TOMATOFS — To control Aphids Vinegar Flies (Drosophila sp.), Anthracnose Septoria Leaf Spot and Early and Late Blight. Do not apply within 3 days of harvest.

CAUTION: Keep Out of Reach of Children

HARMFUL IF SWALLOWED. Avoid breathing of dust. Avoid contact with skin, clothing, and eyes. Avoid contamination of feed and food-stuffs. Wash thoroughly after handling and before eating or smoking. Material may be absorbed through skin.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding on treated areas may be killed. Do not apply when conditions favor drift from the area treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

Do not re-use empty container, destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Buyer assumes all risk of use and/or handling, when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-285

Net Wt. 50 Lbs.



HAVILAND

1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 495.02

ACCEPTED WITH COMMENTS