

# copper maneb fungicide dust 3535

## ACTIVE INGREDIENTS

Copper (from tribasic copper sulfate) . . . . .	3.5%
Maneb (Manganese ethylenebisdithiocarbamate) . . . . .	3.5%
INERT INGREDIENTS . . . . .	93.0%
Total . . . . .	100.0%

## DIRECTIONS

The following directions are furnished by us and are believed to be of value but are in no way guaranteed. Apply at the rate of 50 to 70 lbs. per acre depending on the size and spacing of the plants. For directions as to timing, frequency, and number of applications in your area, consult your State Agricultural Experiment Station or Extension Service. Generally, begin application when disease threatens and repeat at 7 to 10 day intervals or after a rain as necessary.

**CARROTS:** To control Leaf Blight and Cercospora Leaf Spot.

**CELERY:** To control Early <sup>blight</sup> and Late Blight. Remove residues at harvest by washing, stripping and trimming.

**CUCUMBERS, MELON, AND SQUASH:** To control Angular Leaf Spot, Alternaria, Anthracoese, and Downy Mildew.

**TOMATOES, EGGPLANTS, AND PEPPERS:** Angular Leaf Spot, Alternaria, Anthracoese, and Early <sup>blight</sup> and Late Blight. To avoid damage, do not use on tender young plants under glass.

**POTATOES:** To control Early <sup>blight</sup> and Late Blight. Begin at the lower rate and increase rate as vines increase in size.

**ONIONS:** To control Downy Mildew, Botrytis Leaf Blight and Purple Blotch.

**GRAPES:** To control Downy Mildew, Black Rot apply just before bloom, just after, and 10 days later.

## NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.


## CAUTION:

### Keep Out of Reach of Children

**HARMFUL IF SWALLOWED!** May cause irritation of nose, throat, eyes and skin. Do not breathe dust. In case of skin contact wash with water; for eyes, obtain medical attention.

This product is toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from the area treated. Do not contaminate feed or foodstuffs.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.



EPA Reg. No. 595-286



MANUFACTURED BY W. GRAND RAPIDS, MICHIGAN