ACTIVE INGREDIENTS

Parathion (O-O-diethyl O-p-nitrophenyl	
thiophosphate	2.0%
Copper (from tribasic copper sulfate)	
Maneb (Manganese ethylenebisdithiocarbamate).	3.5%
INERT INGREDIENTS:	91.0%
_	100.0%



DANGER: KEEP OUT OF REACH OF CHILDREN

Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Parathion protection. In case of contact, wash immediately with soap and water. Wash with soap and water after handling and before eating or smoking. Wear clean clothing and wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons and animals out of operating areas or vicinity where there may be danger of drift. Workers entering treated fields within 24 hours should wear protective clothing.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

ANTIDOTE

ANTERNAL A CONTRARA OF THE CONTRARA OF A STREET OF A S

FOR TEPNAL CONTRACTOR STATE OF THE STATE OF

DIRECTIONS FOR USE

As recommendations may differ in various agricultural aleas, it is suggested that you contact your State Agricultural Experiment Station or Extension Service for specific rates and timing of applications. If the crop to be dusted is for canning purposes, consult your cannery fieldman before dusting. The following directions are furnished by us and are believed to be of value. This dust should be applied at the rates suggested below depending on the size and spacing of the plants and the method of applications.

CARROTS — To control Aphids, Six Spotted Leafhopper, Alternaria Leaf Blight and Cercospora Blight, apply 25 to 30 lbs. per acre. Begin application when plants are coming up and repeat weekly as long as insects or blight threatens. Do not use on carrots within 15 days of harvest. Do not use treated tops for feeder feed.

CELERY — To control Aphids, Celeryworm and Early and Late Blight, apply 30 to 40 lbs. per acre weekly. Begin dusting as soon as insects or disease threatens. Do not use after celery is half grown or begins to bunch. Remove excess residues by stripping, trimming, and washing.

CUCUMBER, MELON AND SQUASH — To control Cucumber Beetle, Aphids, Stink Bug, Melonworm, Pickleworm, Anthracnose, Cercospora Leaf Spot and Downy Mildew, apply at the rate of 20 to 25 lbs. per acre when insects or disease threatens and repeat at weekly intervals as necessary. Do not use on plants before vining. Do not use within 15 days of harvest on cucumbers or squash nor within 7 days on melons.

ONIONS — To control Downy Mildew, Thrips, Blast (Botrylis Leaf Blight), and Purple Blotch, apply at the rate of 30 to 40 lbs. per acre. Begin dusting 14 days after planting and repeat at 3 to 5 day intervals as needed, if insects appear or blight threatens. Do not apply within 15 days of harvest.

NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. REG. NO. 595-291

NET WT. 50 LBS.



1845 STERLING N.W. GRAND RAPIDS MICHIGAN 49502