diazinon* dust no.2

*Diazinon is a proprietary material of Geigy Agricultural Chemical Co. for the active material O.O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) thiophosphate.

DIRECTIONS

CUCUMBERS, MELONS AND SQUASH — To control Aphids, Leafhoppers, and Miles apply at the rate of 50 to 70 lbs. per acre when insects first appear and repeat as necessary. Do not apply within 7 days of harvest to Summer Squash or Cucumbers, nor within 3 days to Melons or Winter Squash.

ONIONS — To control Onion Thrips apply at the rate of 50 lbs. per acre when insects appear and repeat as necessary. Do not apply within 10 days of harvest.

TOMATOES — To control Aphids, Vinegar Flies (Drosophila sp.) and Dipterous Leafminers apply at the rate of 25 lbs. per acre. Begin when insects first appear and repeat at weekly intervals or as necessary. Do not apply within 3 days of harvest.

CABBAGE, CAULIFLOWER, AND BROCCOLI — To control Aphids, Diamondback Moths, and Imported Cabbageworms apply at the rate of 25 to 50 lbs. per acre. Begin when insects first appear and repeat as necessary. Do not apply within 7 days of harvest on Cabbage or 5 days on Cauliflower or Broccoli.

POTATOES — To control Aphids, Flea Beetles, and Colorado Potato Beetles, apply at the rate of 25 to 40 lbs. per acre. Begin when insects first appear and repeat as needed. Do not apply within 35 days of harvest.

APPLES — To control Codling Moths, Apple Maggots, Aphids, San Jose Scale crawlers, Forbes' Scale Crawlers, and mites apply at the rate of 50 lbs. per acre. Begin at the first cover spray and repeat at weekly intervals or as necessary. Do not apply within 14 days of harvest.

CHERRIES — To control Black Cherry Aphids, Cherry Fruit Flies, and San Jose Scale Crawlers apply at the rate of 25 to 50 lbs. per acre. Begin application at petal fall and continue through cover applications. Do not apply within 10 days of harvest.

PEACHES — To control Aphids, Mites, and San Jose Scale Crawlers apply at the rate of 50 lbs. of dust per acre. Begin at petal fall and continue through cover applications. Do not apply within 20 days of harvest.

STRAWBERRIES — To control Aphids, and Mites apply at the rate of 25 to 50 lbs. per acre. Begin when pests appear and repeat as necessary. Dust should be directed into the crowns of the plants and the plants sufficiently agitated to insure thorough coverage. Do not apply within 5 days of harvest.

COCKROACH CONTROL — (By Pest Control Operators Only) For control of around dwellings and other buildings, around water pipes, cracks and crevices, behind and beneath cabins and areas where roaches crawl or hide. Treatments should be repeated as infestation warrants.

Care should be taken to avoid deposits which could be frequently contacted by children or pets. Do not contaminate food food containers or cooking utensils.

CAUTION:

Keep Out of Reach of Children

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, clothing, and eyes. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling and before eating or smoking. Material may be absorbed through skin.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-292

ACCEPT:
6-20-72 Net Wt. 50 Lbs.
595-292-20 Comment



