

ACTIVE INGREDIENTS

Parathion (O-O-diethyl O-p-nitrophenyl thiophosphate)	2.0%
Copper (from Tri Basic Copper Sulfate)	7.0%
INTERT INGREDIENTS	91.0%



DANGER: Keep Out of Reach of Children

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothes. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Parathion protection. In case of contact, wash immediately with soap and water. Wash with soap and water after handling and before eating or smoking. Wear clean clothing and wash all contaminated clothing with soap and hot water before re-use. Workers entering treated fields within 24 hours should wear protective clothing.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ANTIDOTE

INTERNAL — Give a teaspoon of mustard in a glass of warm water, repeat until vomit is clear. Call a physician immediately! Atropine Sulphate on advice of physician.

EXTERNAL — If skin is contaminated, wash immediately with soap and warm water. If eyes are contaminated, flush with plenty of water and obtain medical attention.

DIRECTIONS FOR USE

As recommendations for the use of Parathion products vary in different agricultural areas, it is suggested that you consult your State Agricultural Experiment Station or Extension Service before using this product. If the crop to be dusted is for canning purposes, consult your cannery fieldman before applying dust. General recommendations for this dust are shown below.

CARROTS — To control aphids, leafhopper and carrot blight, apply dust at the rate of 25 to 35 lbs. per acre depending on the size and spacing of the plants. Begin application early when plants are coming up and repeat applications at weekly intervals as long as insects or blight threatens. Do not apply within 15 days of harvest. For adequate control of leafhoppers, it is necessary to treat margins of fields also. Do not use treated carrot tops for food or feed.

CELERY — To control aphids, early and late blight, apply dust at the rate of 15 to 20 lbs. per acre at weekly intervals. Begin applications as soon as insects appear or blight threatens and repeat at 7 to 10 day intervals as necessary. Do not apply within 21 days of harvest.

CUCUMBERS, MELONS AND SQUASH — To control cucumber beetle, aphids, stink bugs, melonworms and pickleworms, angular leaf spot, alternaria disease, scab and anthracnose, apply 15 to 20 lbs. per acre when insects appear or disease threatens and repeat at 7 to 10 day intervals as necessary. Do not apply before vining or within 15 days of harvest on cucumbers. Do not use within 15 days of harvest on squash or within 7 days of harvest on melons.

POTATOES — To control grasshoppers, aphids and late blight, apply when insects appear or when blight threatens. Use at the rate of 30 to 50 lbs. per acre and repeat at 7 to 10 day intervals as necessary. Do not apply within 5 days of harvest.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. REG. NO. 595-294

ACCEPTED
 5-8-72
 595-294

NET WT. 50 LBS.

UNDER THE FEDERAL INSECT...