RESTRICTED USE PESTICIDE

PARATHION COPPER DUST NO. 27

ACTIVE INGREDIENTS	
Parathion (O O diethyl O p-nitrophenyl thiophosphate)	2.0%
Copper (from Tri Basic Copper Sulfate)	7.0%
INTERT INGREDIENTS	91.0%
	100.0

DANGER: Keep Out of Reach of Children

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothes. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Parathion protection. In case of contact, wash immediately with soap and water. Wash with soap and water after handling and before eating or smoking. Wear clean clothing and wash all contaminated clothing with soap and hot water before re-use. Workers entering treated fields within 24 hours should wear protective clothing.

This product [illegible] fish, birds, and other wildlife. Birds and other wildlife [illegible] treated [illegible] Keep out of any [illegible] [illegible] fee[illegible]

Do [illegible] ing. Stay [illegible]

DIRECTIONS FOR USE

As recommendations for the use of Parathion products vary in different agricultural areas, it is suggested that you consult your State Agricultural Experiment Station or Extension Service before using this product. If the crop to be dusted is for canning purposes, consult your cannery fieldman before applying dust. General recommendations for this dust are shown below.

CARROTS — To control aphids, leafhopper and carrot blight, apply dust at the rate of 25 to 35 lbs. per acre, depending on the size and spacing of the plants. Begin application early when plants are coming up and repeat applications at weekly intervals as long as insects or blight threatens. Do not apply within 15 days of harvest. For adequate control of leafhoppers, it is necessary to treat margins of fields also. Do not use treated carrot tops for food or feed.

CELERY — To control aphids, early and late blight, apply dust at the rate of 15 to 20 lbs. per acre at weekly intervals. Begin applications as soon as insects appear or blight threatens and repeat at 7 to 10 day intervals as necessary. Do not apply within 21 days of harvest.

CUCUMBERS, MELONS AND SQUASH — To control [illegible] the aphids, [illegible] bugs, melonworms and pickleworms, and [illegible] spot, downy mildew, disease, scab and anthra[illegible] apply [illegible] 20 [illegible] [illegible] [illegible] threaten[illegible]

POTATOES [illegible]

NON-WARRANTY

[illegible]

E.P.A. Reg. No. 595-291
E.P.A. Est. No. 10183-MI-4

NET WT. 50 LBS

ACCEPTED
MAY [illegible] 1970

HAVILAND AGRICULTURAL
18 STERLING, N.W. GRAND RAPIDS, MICH. 50[illegible]