

PARATHION 8-E

ACTIVE INGREDIENTS:

Parathion (0,0-diethyl O-p-nitrophenyl phosphorothioate)	80.43%
Petroleum Distillate	8.34%

INERT INGREDIENTS:

Total	<u>11.23%</u>
	100.00%



DANGER: POISON



ANTIDOTE: Internal - Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately! Atropine Sulfate is antidotal. Never give morphine. Never give anything by mouth to an unconscious person. External - If on skin, remove contaminated clothing and immediately wash skin with soap and water; for eyes, flush with plenty of water for at least 15 minutes and call a physician immediately.

DANGER: KEEP OUT OF REACH OF CHILDREN

Poisonous if swallowed, inhaled or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes or on skin or clothing. Wear natural rubber gloves, protective clothing, and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Store material in safe place away from children and away from heat or open flame. DECONTAMINATE AND DESTROY EMPTY CONTAINER.

This product is toxic to fish and wildlife, Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS

Application should be made only by experienced personnel wearing approved safety equipment.

Recommended amount of Haviland Parathion 8-E should be added to nearly filled spray tank and the necessary water added to bring tank to proper level. Agitator should be running during the filling and spraying operation to insure proper distribution of the insecticide. Wettable powders and emulsible liquid should not be added to the spray tank unless previous experience has proved them to be compatible and safe to plants. Rates below are to be used with sufficient water to give thorough coverage.

BEANS: To control Aphids, Armyworms, Leafhopper, Leaf Miner, Leaf Roller and Mexican Bean Beetle, apply 1/2 pt. per acre when insects first appear and repeat application at 7 to 10 day intervals as necessary. Do not apply within 15 days of harvest.

BFETS: To control Aphids, Leafhoppers and Webworm, apply 1/2 pt. per acre as needed. Do not apply within 15 days of harvest (21 days if treated tops are to be used for feed or food).

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS AND TURNIPS:
To control Aphids, Armyworms, Imported Cabbageworm and Thrips, use at the rate of 1/4 to 3/8 pts. per acre when insects or their damage first appear. Do not use within 10 days of harvest on Cabbage and Turnips nor within 7 days of harvest on Cauliflower, Broccoli and Brussels Sprouts.

CORN: To control Armyworms and Corn Rootworm Adults, apply at the rate of 1/4 pt. per acre when insects appear. To control European Corn Borer, rate should be increased to 1/2 pt. per acre. Do not apply within 12 days of picking or cutting for forage.

CUCUMBERS, MELONS AND SQUASH: To control Aphids, Cucumber Beetles, Melonworms and Pickleworms, apply at the rate of 1/4 to 1/2 pt. per acre. Do not apply when plants are wet or before runners begin to form. Do not apply within 15 days of harvest on cucumbers and squash or within 7 days of harvest on melons.

LETTUCE: To control Aphids and Leafhoppers, apply at the rate of 1/2 pt. per acre. Application should not be made within 21 days of harvest on leaf and bib lettuce or within 7 days of harvest on head lettuce.

PEAS: To control Aphids and Pea Weevil, apply at the rate of 1/2 pt. per acre. Do not apply within 10 days of harvest.

PEPPERS AND TOMATOES: To control Aphids, Flea Beetles and Leafhoppers, apply at the rate of 1/4 pt. per acre. Do not apply within 10 days of any picking on tomatoes or within 15 days of any picking on peppers.

POTATOES: To control Aphids, Flea Beetles and Leafhoppers, apply at the rate of 1/4 pt. per acre. Do not apply within 5 days of harvest.

SUGAR BEETS: To control Aphids, Blister Beetles and Leafhoppers, apply at the rate of 1/2 pt. per acre. Do not make application within 15 days of harvest.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-302

Net Contents _____ GAL.

HAVILAND AGRICULTURAL CHEMICAL COMPANY

GRAND RAPIDS, MICHIGAN

1845 STERLING, N. W.