


ZINC PHOSPHIDE DEMOUSER

ACTIVE INGREDIENTS

Zinc Phosphide.....94%

INERT INGREDIENTS.....6%

Total.....100%



*(Handwritten)*

POISON



POISON

*(Handwritten)*

DANGER- Keep Out Of Reach Of Children

"SEE SIDE PANEL FOR ADDITIONAL WARNINGS AND ANTIDOTE STATEMENT"

Net Weight 2 Lbs.

HAVILAND AGRICULTURAL CHEMICAL CO., 1845 Sterling, N.W., Grand Rapids, Michigan

(Center Panel)

Suggested Bait Formula:

- 100 Pounds of Cracked Corn
- 2 Pounds of Zinc Phosphide
- 1 Quart Vegetable Oil
- 1/2 Ounce Methyl Green Dye

DIRECTIONS FOR MAKING FORMULA: The cracked corn may be dyed by dissolving 1/2 ounce Methyl Green in one gallon water and pouring it over corn in a large tub. Mix thoroughly. Allow to dry thoroughly. Alternate method: Mix the dye with the dry Zinc Phosphide before adding to the oil. The prepared grain will then turn green with contact of moisture from the ground, dew, frost or rain.

Mix the Zinc Phosphide (or Zinc Phosphide and dye) with the oil to form a thin paste. Pour paste over grain in large tub. Use hoe to mix the mass very completely so the grains are evenly coated. More poison will stick to the broken corn surfaces than to the unbroken surfaces. For mixing large quantities, any drum type mixer may be used. Thoroughly scrub out with soap and water after using.

(Left Side Panel)

DIRECTIONS FOR STORING: Zinc Phosphide bait is poisonous to humans and animals. The material must be stored safely. Do not store bait in tightly closed buildings where animals or people are present. Keep dry. Label all containers "POISON" and use them for no other purpose.

DIRECTIONS FOR USE:

SPOT OR TRAIL BAITING: Place one teaspoon of bait in surface trails or at the mouth of holes leading to underground burrow systems. Two to four bait spots should be made near the base of each infested tree. Cover bait by pulling overhanging grass into place. Do not disturb the runway-system. Baiting should be conducted at the rate of 2 to 3 pounds per acre.

BROADCAST BAITING: May be applied by cyclone seeder, or by hand. When applying by hand, throw tablespoon amounts into heavy cover along tree rows, rock outcrops, and fence lines. Apply at the rate of 6 to 10 pounds per acre. Do not broadcast over growing crops.

MACHINE BAITING: Use a modified corn planter with plates set to drop one kernel per inch along a line just inside the drip line on both sides of the tree. Bait should be applied at the rate of 3 to 6 pounds per acre.

ZINC PHOSPHIDE DEMOUSER

SPECIAL CAUTIONS: Do not allow bait to be exposed on bare ground. Store poisoned grains in location not accessible to children, pets, poultry, and domestic animals.

WARNING: POISONOUS IF SWALLOWED! Do not breathe dust or vapors. Avoid contact with skin. Wash thoroughly after handling and before eating or smoking. Store away from and do not contaminate feed or food-stuffs. Keep away from children, pets, and domestic animals, and keep them out of baited areas. Birds feeding in treated areas may be harmed.

Toxic to wildlife.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ANTIDOTE: Give 1 tablespoon of mustard in a glass of warm water, and repeat until vomit fluid is clear. Avoid the use of oils. Have patient lie down and keep warm. Call a physician immediately.

USDA Reg. No. 595-305

(Right Side Panel)

HAVILAND AGRICULTURAL CHEMICAL CO.  
1845 Sterling, N. W.  
Grand Rapids, Michigan 49502