

chlordane emulsifiable concentrate 8-E

(Contains 8 lbs. active per gallon)

ACTIVE INGREDIENTS:

*Chlordane Technical	72.0%
Aromatic Petroleum Distillate	23.0%
INERT INGREDIENTS	5.0%

Total 100.0%

*Equivalent to 43.2% Octachloro-4, 7-methanotetrahydroindane and 28.8% related compounds.

For specific applications, follow the general directions below or consult your local state agricultural experimental station for specific recommendations in the area in which the material is to be used.

WARNING: Keep Out of Reach of Children

Harmful if swallowed. Contact with skin causes toxic symptoms. Do not apply to forage crops that are to be fed to dairy animals or to animals being finished for slaughter. Avoid inhalation and contamination of feed and foodstuffs. Do not apply to any crop after edible parts begin to form. Do not apply to food crops within 30 days of harvest. Avoid contact with skin and eyes. For contact with skin, wash thoroughly with soap and water; for eyes, flush with plenty of water and obtain medical attention. If taken internally, induce vomiting and call a physician.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS

CROP USE: CABBAGE, CAULIFLOWER, BROCCOLI (transplanting time only)
To control Cabbage Maggot use ½ pt. of material to 100 gallons of water and use 1/3 pt. of this mixture to each plant as a transplant water treatment when plants are being set in the field. Also apply treatment to the soil surface 10 days after plants are first treated. A directed spray should be used so as not to spray newly set plants. Do not apply treatment after edible parts begin to form. Do not apply more than 1 qt. of concentrate per acre (2 lbs. actual Chlordane).

POTATOES, CELERY, PARSNIPS – To control Wireworms apply 2½ pts. to 5 qts. per acre in sufficient amount of water to insure good distribution as a soil treatment prior to planting. Use the lower rate for the light mineral soils and the higher rate for organic or muck soils.

CORN – To control Corn Rootworm apply 3 qts. of material in sufficient water to cover each acre. Apply as a pre plant soil treatment and disc into the upper 3 or 4 inches of soil.

STRAWBERRIES – To control Strawberry Root Weevil Larvae apply at the rate of 5 qts. per acre and disc into the upper 6 inches of soil. Application should be made in the spring before planting.

DRY ONIONS (only) – To control Onion Thrips and Onion Maggot apply 1 pt. of material per 100 gallons of water and apply at the rate of 50 to 100 gallons per acre beginning when the plants are six inches high and repeat at weekly intervals. Use the lower rate when the plants are small and increase rate as size of plants increase. Do not apply this product to green or spring onions.

NON-CROP USE

LAWN & ORNAMENTAL TURF OR SOD AREAS, ROADWAYS, DITCH BANKS – to control White Grubs and Japanese Beetle Larvae dilute ¼ pt. of material in sufficient water to cover 1,000 sq. ft. or use 1½ gallons of actual Chlordane 8-E per acre. Spray area to be treated and then thoroughly wet with additional water to insure placing the toxicant into position of maximum effectiveness. To control Ants and Wireworms use 1 to 1½ pts. per acre (4 to 6 ounces per 10,000 sq. ft.) in sufficient water to insure adequate coverage and follow with additional water to place toxicant into upper layer of soil. To control Chiggers and Lawn Moths (Sod Web-worms) apply at the rate of ½ pt. per 4,000 sq. ft. or 2 qts. per acre. Make application after first thoroughly wetting area to be treated and do not follow application with additional water for at least 2 or 3 days. To control Mosquitoes prepare spray by diluting 1 pt. of material in 100 gallons of water. Apply 20 to 40 gals. per acre. In spraying, be sure to pay attention to tall grasses around buildings and shrubs. Do not apply spray to large trees or shrubs where birds are roosting or nesting. Do not apply spray to food crops. Do not graze livestock on treated areas. Keep children and pets off treated areas until the areas have been well watered and dried completely.

SUBTERRANEAN TERMITES - Directions for Non-Professional Use

Use 1% Chlordane 8E in a 99% water emulsion as follows:

Dilute in ratio of 1 gallon of Chlordane Emulsifiable concentrate 8E to 99 gallons of water.

Buildings Having Basements or Crawl Spaces Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 595-321

ACCEPTED
8-21-72
595-321

NET CONTENTS 5 GALLONS



HAYILAND

DIRECTIONS

Feed at the rate of 1.5 pounds per head per day.
Maintain a supply of clean water near feeding areas.

WARNING

DO NOT FEED TO LACTATING DAIRY ANIMALS. OCCASIONALLY, AN ANIMAL MAY BE SENSITIVE TO PHENOTHIAZINE RESULTING IN SYMPTOMS SUCH AS TEMPORARY BLINDNESS AND LOSS OF SKIN OR HAIR. IF TOXIC SYMPTOMS DEVELOPE, TAKE ANIMALS OFF DRUG AND AVOID EXPOSURE TO SUNLIGHT FOR SEVERAL DAYS. DO NOT ADMINISTER TO ANIMALS SHOWING SYMPTOMS OF ACUTE RESPIRATORY CONDITIONS.

NOTE

This product kills horn fly and face fly maggots in manure but does not control existing adult flies. If necessary, spray cattle for maximum protection. Because face flies migrate from farm to farm, it is essential that a larvicide program be employed on an area basis to assist in reducing the adult face fly population.

A-3234