

NET CONTENTS:

HOW TO USE PURINA POULTRY INSECTICIDE

AS A ROOST PAINT

To help control ~~Lice and Mites~~—Brush Poultry Insecticide on top of cleaned perches or roost poles before birds go to roost. Apply 1 pint per 150 feet. Fumes rise and kill lice and mites in feathers. If necessary, repeat in 8 to 10 days to kill lice that may hatch later.

FOR CAGED LAYERS

To help control Lice and Mites—Apply Poultry Insecticide with a paint roller or brush on the underside of floor wires of cages. Allow 3 to 5 days for the treatment to be fully effective. Repeat in 30 days if birds become reinfested.

CAUTION: Collect eggs before treatment to prevent possible contamination. Do not apply Poultry Insecticide to the wires of egg collection racks.

AS A SPRAY FOR CHICKEN HOUSES

To help control Lice, Mites, Bedbugs, and Flies—Mix 1 quart of Poultry Insecticide with 3 quarts of water. Spray walls, ceilings, roosts, nest boxes and dropping boards; apply 1 gallon of spray per 500 square feet. Provide clean nesting material after spraying.

FOR DIRECT APPLICATION ON BIRDS

To help control Lice and Mites—As a spray: Mix 1 pint of Poultry Insecticide with enough water to make 1 gallon of spray solution. Spray birds at the rate of 1 gallon of mixed solution to 100-150 birds depending on size. Treat again when necessary.

To help control Northern Fowl Mites, Chicken Body Lice, and Shaft Lice—For tail dipping: Mix ½ pint of Poultry Insecticide with enough water to make 1 gallon of dipping solution. Hold bird by wings; dip 3 or 4 inches of tail with enough force to open the feathers; thoroughly wet the vent and surrounding areas. Treat again in seven to ten days as needed.

CAUTION: Vapor harmful. Harmful or fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Atropine and 2-PAM are antidotal.

CAUTION: Avoid contact with skin, eyes and clothing. Avoid breathing spray mist. Always use freshly mixed solutions. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs. If product has been stored under cold conditions, shake before using.


This product is toxic to fish. Keep out of any body of water. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in safe place.

705 B


D-7263

Printed in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH



PURINA
POULTRY
INSECTICIDE
(ROOST PAINT)



Like
Active Ingredients
*Malathion . . .
Xylene . . .
Petroleum Oil
Inert Ingredients

*O,O-dimethyl dit

**CAUTION:
SEE BACK**

RALSTON
GENERAL OFFICES •

A POULTRY INSECTICIDE

ROOST PAINT

Poultry Insecticide on top of cleaned perches or
by 1 pint per 150 feet. Fumes rise and kill lice and
in 8 to 10 days to kill lice that may hatch later.

AGED LAYERS

Poultry Insecticide with a paint roller or brush on
new 3 to 5 days for the treatment to be fully effec-
tively infested.

To prevent possible contamination. Do not apply
to collection racks.

FOR CHICKEN HOUSES

Fleas—Mix 1 quart of Poultry Insecticide with
roosts, nest boxes and dropping boards; apply
Provide clean nesting material after spraying.

APPLICATION ON BIRDS

Mix 1 pint of Poultry Insecticide with enough
water. Spray birds at the rate of 1 gallon of mixed
solution. Treat again when necessary.

Body Lice, and Shaft Lice—For tail dipping: Mix
with enough water to make 1 gallon of dipping solution.
Dip tail with enough force to open the feathers;
repeat in areas. Treat again in seven to ten days as needed.

Fatal if swallowed. If swallowed, induce vomiting.
Activated 2-PAM are antidotal.


Wash hands and clothing. Avoid breathing spray mist.
Wash thoroughly with soap and water after using.
Do not use on stuffed birds. If product has been stored under cold

any body of water. Do not reuse empty container.
Recycle or discard in safe place.

63


Printed in U.S.A.

BY PURINA RESEARCH



NET CONTENTS:


32 FL. OZ. (1 QT.)



PURINA®

POULTRY INSECTICIDE (ROOST PAINT)

U.S.D.A. REG. NO. 602-36



Helps in Controlling
Lice—Mites—Bedbugs—Fleas

Active Ingredients:

*Malathion	4.00%	(.28 lbs/gal)
Xylene	25.00%	(1.77 lbs/gal)
Petroleum Oil	65.75%	(4.66 lbs/gal)
Inert Ingredients	5.25%	
	<u>100.00%</u>	

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

MANUFACTURED BY

RALSTON PURINA COMPANY
GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MO. 63199