

Reason For Issue:
Changes per EPA. For Submission to EPA.

Date Issued: 1/17/90 (Green) 602-36 1/3
P.M. 16 1-25-90
Prev. Date Issued: 4/15/88

Front Panel

(9-Square Logo)

Contains
CYTHION
The Premium Grade
Malathion
Registered Trademark of
American Cyanamid
Company

PURINA
BRAND

POULTRY INSECTICIDE
(ROOST PAINT)

EPA Reg. No. 602-36

Helps in controlling
Lice - Mites - Bedbugs - Fleas

(3-Row Checkerband)

ACTIVE INGREDIENTS:

Malathion: 0,0-dimethyl phosphorodithioate of diethyl
mercaptosuccinate..... 4.00% (.28 lbs/gal)

INERT INGREDIENTS..... 96.00%
100.00%

Inert ingredients include Aliphatic petroleum hydrocarbons and Xylene

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

PROTECT FROM FREEZING

NET 32 FL OZ (1 QT)


D-7263

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812
St. Louis, Missouri 63166-6812 U.S.A.

(NOTE: Type size and prominence
will conform with requirements
for precautionary labeling.)

NET 32 FL OZ
PURINA POULTRY
BRAND
INSECTICIDE

Not Reviewed. Registered in accordance with
Regulation. Not reviewed for Phase 2
Response. Agency to alert Change in Status.



Back Panel
HOW TO USE PURINA POULTRY INSECTICIDE
BRAND

AS A ROOST PAINT

To help control Lice and Mites- Brush Poultry Insecticide on top of cleaned perches or roost poles before birds go to roost. Apply 1 pint per 150 feet. Fumes rise and kill lice and mites in feathers. If necessary, repeat in 8 to 10 days to kill lice that may hatch later.

FOR CAGED LAYERS

To help control Lice and Mites - Apply Poultry Insecticide with a paint roller or bush on the underside of floor wires of cages. Allow 3 to 5 days for the treatment to be fully effective. Repeat in 30 days if birds become reinfested.

CAUTION: Collect eggs before treatment to prevent possible contamination. Do not apply Poultry Insecticide to the wires of egg collection racks.

AS A SPRAY FOR CHICKEN HOUSES

To help control Lice, Mites, Bedbugs, and Fleas - Mix 1 quart of Poultry Insecticide with 3 quarts of water. Spray walls, ceilings, roosts, nest boxes and dropping boards; apply 1 gallon of spray per 500 square feet. Provide clean nesting material after spraying.

FOR DIRECT APPLICATION ON BIRDS

To help control Lice and Mites - As a spray: Mix 1 pint of Poultry Insecticide with enough water to make 1 gallon of spray solution. Spray birds at the rate of 1 gallon of mixed solution to 100 - 150 birds depending on size. Treat again when necessary.

To help control Northern Fowl Mites, Chicken Body Lice, and Shaft Lice - For tail dipping: Mix 1/2 pint of Poultry Insecticide with enough water to make 1 gallon of dipping solution. Hold bird by wings; dip 3 or 4 inches of tail with enough force to open the feathers; thoroughly wet the vent and surrounding areas. Treat again in seven to ten days as needed.

CAUTION: Vapor harmful. Harmful or fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Atropine and 2-PAM are antidotal.

CAUTION: Avoid contact with skin, eyes and clothing. Avoid breathing spray mist. Always use freshly mixed solutions. Wash thoroughly with soap and water after using.

Avoid contamination of feed and foodstuffs. If product has been stored under cold conditions, shake before using.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces. Do not allow contact with treated surface until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in cool, dry area out of reach of children and animals.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.

QUALITY CONTROLLED BY PURINA RESEARCH

EPA Est. No. 602-MO-1

8901 J

Printed in U.S.A.

