

NE

HOW TO USE PURINA LIQUID STOCK SPRAY

MIXING DIRECTIONS

Beef Cattle and Hogs: For the control of Horn Flies, Lone Star Ticks, Winter Ticks, Gulf Coast Ticks, Spinose Ear Ticks, Lice and Sarcoptic Mange Mites—Mix 1 gallon of Liquid Stock Spray in 100 gallons of water (2 1/2 tablespoonfuls to 1 gallon of water) for either spraying or dipping.

Sheep and Goats: For the control of Sheep Ticks (Keds) and Lice—Mix 1 quart of Liquid Stock Spray in 125 gallons of water for spray solution or in 250 gallons of water for dipping vat solution.

USE DIRECTIONS:

Mix spray or dipping vat solutions thoroughly before using and keep well mixed while spraying or dipping animals. Refill dipping vats at indicated rates when fresh water must be added. Clean out entire vat when solution becomes filthy. Repeat treatment at monthly intervals, if necessary. *Do not* spray animals confined in closed buildings. *Do not* spray horses or mules. Avoid treating animals in cold, stormy weather.

Cattle: Apply up to 1 gallon of spray solution per animal, except calves less than 6 months old. Do not directly spray or dip calves less than 6 months old. For Gulf Coast Ticks and Spinose Ear Ticks, apply at low pressure to each ear and to head, neck and shoulder region.

Sheep and Goats: Apply 2 to 4 quarts of spray solution per animal at low pressure to lambs and adults after spring shearing. Use high pressure to penetrate fleece

during fall treatment. Hold animals in dipping vats for 1/2 minute and duck head twice for an instant while swimming through vat.

Hogs: Spray or dip entire animal so as to thoroughly wet the skin, including the underside. Do not treat sows within 2 weeks before farrowing nor for at least 3 weeks after farrowing.

WARNING: May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately.

Do not breathe spray mist. Do not get in eyes, on skin or on clothing. May be absorbed through skin. Wash face and hands thoroughly. Avoid contamination of feed, foodstuffs, feeders, waterers and food equipment.

Do not use on dairy animals or in dairy barns. Do not treat lambs, kids or pigs that are in the suckling stage. Do not treat weak or emaciated animals. Do not spray horses or mules. Do not dip animals within 60 days of slaughter. Do not spray animals within 30 days of slaughter. Do not treat animals under 3 months of age. Dip only healthy adult animals. Avoid treating overheated or sick animals.


Toxic to fish and wildlife. Do not contaminate streams or ponds with spray or dipping vat solution. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

D-8975

Printed in U.S.A.




QUALITY CONTROLLED BY PURINA RESEARCH

ACCEPTED
 JAN 18 1973
 UNDER THE FEDERAL INSECTICIDE
 ACT AND RODENTIDE ACT
 FOR FILING RECORDS OF REGISTERED
 TO ALL GOVERNMENTS


NET CONTENTS: 1 GALLON

EPA REG. NO. 602-37-AA

PURINA
LIQUID



STOCK SPRAY



FOR SPRAYING OR DIPPING VATS

KILLS... HORN FLIES, TICKS, LICE, FLEAS AND SARCOPTIC MANGE MITES

Contains 50% Toxaphene and 3% Lindane (by volume)

Active Ingredients:

*Toxaphene	45.0%
Gamma Isomer of Benzene Hexachloride, (from Lindane)	2.7%
Aromatic Petroleum Derivative Solvents	20.0%
Petroleum Hydrocarbons	19.3%
Xylene	5.0%
Inert Ingredients	8.0%

*Technical chlorinated camphene (chlorine content 67-69%) 100.0%

WARNING - KEEP OUT OF REACH OF CHILDREN

SEE OTHER WARNINGS ON BACK PANEL

MANUFACTURED BY

RALSTON PURINA COMPANY

GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63188

