

HOW TO USE PURINA MANGE CONTROL

Purina Mange Control can be mixed with water to make a spray or dip for hogs, beef cattle, sheep, non-milking goats, or dogs. Mange Control can also be mixed with oil to make a medicated oil for hogs. All animals must be treated thoroughly, including inside of ears. Two applications, a week apart, may be necessary. Infested premises should be sprayed using 1 pint of Mange Control to 16 gallons of water; replace litter.

FOR WATER SPRAY OR DIP

Mixing Directions—Mix 1 pint of Mange Control with 16 gallons of water, or 1 ounce (2 tablespoons) per gallon. (See special instructions for calves, other young animals and dogs.)

HOW TO USE—For Mange or Scabies

Hogs—Use 2 to 3 quarts of mixed solution per animal.

Beef Cattle—Use 2 quarts of mixed solution per adult animal.

For Calves and other young animals (beyond the minimum ages as listed under warning) use ½ pint of Mange Control to 16 gallons of water (1 tablespoon per gallon).

Sheep and Non-Milking Goats—Use 1 to 2 gallons of mixed solution per adult animal. If fleece is thick, spray pressure of 350 to 400 lbs. may be necessary.

Dogs—Use ½ pint of Mange Control to 16 gallons of water (1 tablespoon per gallon). Bathe or dip. Repeat treatment weekly if reinfestation occurs.

MEDICATED OIL SOLUTION FOR HOGS

Mixing Directions—Mix 1 pint of Mange Control with 16 gallons of stove oil or diesel fuel oil, or 1 ounce (2 tablespoons) per gallon.

HOW TO USE—(For Hogs Only)

Hand Application or Spraying—Use sprinkling can or sprayer to apply an even coat of oil over the entire surface of head and body. Treat inner ear surfaces and underside of body, especially between thighs, with brush or swab.

Hog Oilers and Rubbing Posts—Keep oilers filled and rubbing posts saturated with medicated oil solution as a preventive measure. If outbreaks occur, use hand application or sprayer to insure complete coverage.

WARNING—Do not treat animals within 30 days prior to giving birth to young. Do not treat nursing animals. Do not treat pigs less than 3 weeks old, calves less than 3 months old, or dogs less than 4 months old. Do not treat animals within 60 days of slaughter. Keep oiled hogs out of direct sunlight for 24 hours. Do not use on dairy animals.

CAUTION—Avoid contact with skin, eyes, or clothing. Avoid contamination of feed and foodstuffs. Avoid breathing of vapor or mist. May be absorbed through skin. Wash hands and face thoroughly.

DANGER—Harmful or fatal if swallowed. If swallowed induce vomiting. Call physician immediately.

6512D

D-5972

PROTECT FROM FREEZING

Printed in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH

ACCEPTED WITH COMMENTS


NET CONTENTS

1 LIQUID PINT

PURINA[®]
MANGE CONTROL



U.S.D.A. REG. NO. 602-39



TO BE MIXED WITH WATER OR OIL

Controls . . . Mange (Sarcoptic) and Scabies (Psoroptic) of Farm Animals

Mange (Sarcoptic), Ticks, Fleas and Lice on Dogs

Active Ingredients:

Gamma Isomer of Benzene Hexachloride (from Lindane)

Aromatic Petroleum Derivative Solvent

Inert Ingredients

6.5%
85.5%
8.0%

100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

Manufactured by

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Mo. 63199

HOW TO USE PURINA MANGE CONTROL

Purina Mange Control can be mixed with water to make a spray or dip for hogs, Beef cattle, sheep, non-milking goats, or dogs. Mange Control can also be mixed with oil to make a medicated oil for hogs. All animals must be treated thoroughly, including inside of ears. Two applications, a week apart, may be necessary. Infested premises should be sprayed using 1 pint of Mange Control to 16 gallons of water; replace litter.

FOR WATER SPRAY OR DIP

Mixing Directions Mix 1 pint of Mange Control with 16 gallons of water, or 1 ounce (2 tablespoons) per gallon. See special instructions for calves, other young animals and dogs.

HOW TO USE—For Mange or Scabies

Hogs Use 2 to 3 quarts of mixed solution per animal.

Beef Cattle Use 2 quarts of mixed solution per adult animal.

For Calves and other young animals, beyond the minimum ages as listed under warning, use 1/2 pint of Mange Control to 16 gallons of water (1 tablespoon per gallon).

Sheep and Non-Milking Goats Use 1 to 2 gallons of mixed solution per adult animal. If fleece is thick, spray pressure of 350 to 400 lbs. may be necessary.

Dogs Use 1/2 pint of Mange Control to 16 gallons of water (1 tablespoon per gallon). Bathe or dip. Repeat treatment weekly if reinfestation occurs.

MEDICATED OIL SOLUTION FOR HOGS

Mixing Directions Mix 1 pint of Mange Control with 16 gallons of stove oil or diesel kerosene, or 1 ounce (2 tablespoons) per gallon.

HOW TO USE—(For Hogs Only)

Hand Application or Spraying Use sprinkling can or sprayer to apply an even coat of oil over the entire surface of head and body. Treat inner ear surfaces and underside of body, especially between thighs, with brush or swab.

Hog Oilers and Rubbing Posts Keep oilers filled and rubbing posts saturated with medicated oil solution as a preventive measure. If outbreaks occur, use hand application or sprayer to insure complete coverage.

WARNING Do not treat animals within 30 days prior to giving birth to young. Do not treat nursing animals. Do not treat pigs less than 3 weeks old, calves less than 3 months old, or dogs less than 4 months old. Do not treat animals within 60 days of slaughter. Keep oiled hogs out of direct sunlight for 24 hours. Do not use on dairy animals.

CAUTION Avoid contact with skin, eyes, or clothing. Avoid contamination of feed and foodstuffs. Avoid breathing of vapor or mist. May be absorbed through skin. Wash hands and face thoroughly.

DANGER Harmful or fatal if swallowed. If swallowed induce vomiting. Call physician immediately.

4512D

D-5972

PROTECT FROM FREEZING

Made in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH

NET CONTENTS

1 LIQUID PINT

PURINA 
MANGE CONTROL

U.S.D.A. REG. NO. 602-39

TO BE MIXED WITH WATER OR OIL

Controls . . . Mange (Sarcoptic) and Scabies (Psoroptic) of Farm Animals
Mange (Sarcoptic), Ticks, Fleas and Lice on Dogs

Active Ingredients:

Gamma Isomer of Benzene Hexachloride (from Lindane)

Aromatic Petroleum Derivative Solvent

Inert Ingredients

6.5%

85.5%

8.0%

100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

Manufactured by

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Mo. 63199