

CONTENTS: 1 LIQUID QUART

U.S.D.A. REG. NO. 602-46

NA MALATHION SPRAY

application. For the protection of stored grains such as wheat, oats, rice, corn, rye, barley, grain sorghum, and field of garden seeds, against confused flour beetle, rice weevil, granary weevil, saw-tooth grain beetle, flat grain beetle, red flour beetle, rusty grain beetle, lesser grain borer, and Indian meal moth—apply 1 pint of Malathion Spray in 2-5 gallons of water per 1,000 bushels. Apply as the grain is being loaded or turned into final storage.

Home Gardens: Mix 2 teaspoonfuls to 1 gallon of water (1/2 pint in 25 gallons of water). Stir and keep uniformly mixed while spraying. Spray plants thoroughly and repeat as necessary. Do not spray within 7 days of harvest or when plants are in bloom. Do not apply to pecans within 60 days of harvest. **Vegetables**—Aphids (Plant Lice), Spider Mites, and Leafhoppers on Tomatoes, Peas, Beans, Potatoes, Cabbage, Cauliflower, Mustard, Cucumbers, Squash, and Melons; Mexican Bean Beetle on Beans; Imported Cabbage Worm and Cabbage Looper on Cabbage or Cauliflower. **Fruits and Nuts**—Aphids, Mites, Red-banded leaf Roller, Bud Moth, Codling Moth, Plum Curculio, and Mealybug on Apple, Peach, Cherry, Plum, Pear, Pecan and Grape. **Ornamental Shrubs**—Aphids, Spider Mites, Rose Leafhopper, Bagworms, Tent Caterpillars, Birch Leafminer, and Scurfy Scale. Do not spray ferns and crassula.

CAUTION: Avoid breathing spray mist. Avoid prolonged contact with skin, eyes, and clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs. Avoid contamination of milk, milk equipment, and water. Do not use where milk is stored or processed.

Do not spray curcubits unless plants are dry. May cause injury to McIntosh and Cortland apples, Bosc pears, Ribier grapes, and sweet cherries.

DANGER: Vapor harmful. Harmful or fatal if swallowed. If swallowed do not induce vomiting. Call physician immediately.


Printed in U.S.A.

D-5834

ED BY PURINA RESEARCH

PURINA[®] MALATHION SPRAY

PREMIUM
GRADE



FOR INSECT CONTROL: FARM BUILDINGS- POULTRY-LIVESTOCK-STORED GRAINS-GARDENS

Active Ingredients:	
*Malathion.....	50.0%
Xylene.....	41.8%
Inert Ingredients.....	8.2%
	<hr/>
	100.0%

(*)O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS ON BACK PANEL**

Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Square • St. Louis, Missouri