

## HOW TO USE PURINA FLY BAIT

**Dry Bait:** Remove lid and punch in perforated holes on inner liner. Replace lid when not in use.

**Syrup Paint:** Dissolve contents of Fly Bait container (1 pound) in 1/2 pint (one cupful) of water. Use brush to paint on surfaces where flies gather.

### DIRECTIONS FOR USE

**Dairy Barns, Loafing Sheds, Animal Shelters, Stables, Outdoor Areas:** Scatter Fly Bait on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed rooms, alleyways and window sills. On freshly limed floors scatter Fly Bait on clean feed sacks or on paper.

Or brush Fly Bait dissolved in water on supporting posts, rafters, along ceiling cracks and on outer surfaces of animal enclosures. Keep out of reach of animals. Repeat procedure as necessary for practical fly control.

**Caged Poultry and Turkeys on Wire:** Scatter Fly Bait around doorways and along edge of dropping area. Or brush Fly Bait dissolved in water on cage supports, along edges of rafters and around doorways. Repeat as necessary.

**Dog Kennels and Furbearing Animal Units:** Scatter Fly Bait outside dog runways, on window sills, ledges and other surfaces out of reach of the animals. For cage units apply along edges of dropping area. Or brush Fly Bait dissolved in water on cage supports and other surfaces where flies gather. Do not apply to surfaces within reach of animals. Repeat as necessary.

**WARNING:** Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly after applying. Do not contaminate feed and foodstuffs. Keep out of reach of children. Do not apply where poultry or animals, especially young calves, can lick it. Do not use in homes or where milk is processed or stored. Atropine is antidotal.

PAT. NO. 3,160,556

666C

D-3763

PRINTED IN U.S.A.

NET WEIGHT

1 POUND AVDP.



**PURINA<sup>®</sup>**  
**FLY BAIT**



602-53

**(SUGAR BASE)**

U.S.D.A. REG. NO. 602-53



**CONTAINS DDVP & DIPTEREX\*\***

**Use as Dry Bait or Syrup Paint**

<b>Active Ingredients:</b>	
2,2-Dichlorovinyl dimethyl phosphite*.....	0.23%
Related Compounds.....	0.02%
**Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate.....	0.25%
<b>Inert Ingredients.....</b>	<b>99.50%</b>
	<b>100.00%</b>

\*DDVP

\*\*Reg. U.S. Pat. Off. by Farbenfabriken Bayer, A. G. Chemagro Corporation Licensee. U.S. Patent No. 2,701,225

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**SEE OTHER WARNINGS ON BACK PANEL**

Manufactured By

**RALSTON PURINA COMPANY**

GENERAL OFFICES • CHECKERBOARD SQUARE

## HOW TO USE PURINA FLY BAIT

**Dry Bait:** Remove lid and punch in perforated holes on inner liner. Replace lid when not in use.

**Syrup Paint:** Dissolve contents of Fly Bait container (1 pound) in 1/2 pint (one cupful) of water. Use brush to paint on surfaces where flies gather.

### DIRECTIONS FOR USE

**Dairy Barns, Loafing Sheds, Animal Shelters, Stables, Outdoor Areas:** Scatter Fly Bait on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed rooms, alleyways and window sills. On freshly limed floors scatter Fly Bait on clean feed sacks or on paper.

Or brush Fly Bait dissolved in water on supporting posts, rafters, along ceiling cracks and on outer surfaces of animal enclosures. Keep out of reach of animals. Repeat procedure as necessary for practical fly control.

**Caged Poultry and Turkeys on Wire:** Scatter Fly Bait around doorways and along edge of dropping area. Or brush Fly Bait dissolved in water on cage supports, along edges of rafters and around doorways. Repeat as necessary.

**Dog Kennels and Furbearing Animal Units:** Scatter Fly Bait outside dog runways, on window sills, ledges and other surfaces out of reach of the animals. For cage units apply along edges of dropping area. Or brush Fly Bait dissolved in water on cage supports and other surfaces where flies gather. Do not apply to surfaces within reach of animals. Repeat as necessary.

**WARNING:** Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly after applying. Do not contaminate feed and foodstuffs. Keep out of reach of children. Do not apply where poultry or animals, especially young calves, can lick it. Do not use in homes or where milk is processed or stored. Atropine is antidotal.

PAT. NO. 3,160,556

6770

D-3763

PRINTED IN U.S.A.

NET WEIGHT

1 POUND AVDP.

# PURINA<sup>®</sup> FLY BAIT



(SUGAR BASE)

U.S.D.A. REG. NO. 602-53

CONTAINS DDVP & DIPTEREX\*\*

## Use as Dry Bait or Syrup Paint

Active Ingredients:	
2,2-Dichlorovinyl dimethyl phosphlate*	0.23%
Related Compounds	0.02%
**Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate	0.25%
Inert Ingredients	99.50%
	100.00%

\*DDVP

\*Reg. U.S. Pat. Off. by Farbenfabriken Bayer, A. G., Clematis Corporation Licensee, U.S. Patent No. 2,791,225

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

SEE OTHER WARNINGS ON BACK PANEL

Manufactured By

**RALSTON PURINA COMPANY**

GENERAL OFFICES • CHECKERBOARD SQUARE