

**HOW TO USE
PURINA POULTRY
DUSTING POWDER**

AS A LITTER TREATMENT

To Control Northern Fowl Mites, Chicken Mites, Lice, Fleas, and Bedbugs: Apply Poultry Dusting Powder to the litter at the rate of 1 pound to 40 square feet with a hand or power duster. For small areas, use a scoop or scatter by hand. Distribute dust evenly. Also dust roosts and adjacent areas. Avoid direct contamination of eggs. Do not treat nest litter. Repeat as necessary.

FOR DIRECT APPLICATION TO BIRDS

To Control Northern Fowl Mites, Chicken Mites, Lice and Fleas: **Individual treatment:** Shake Poultry Dusting Powder into the feathers, particularly around the vent, neck, and breast regions. Ruffle the feathers so that the powder can reach the skin. Use 1 pound per 100 birds. Repeat dusting in 4 weeks, if necessary. Hand application to birds for chicken mite and fleas is a supplement to litter treatment for control of these pests.

Caged Poultry: Use Poultry Dusting Powder in a hand duster. Hold nozzle so it touches the feathers and apply one puff to both the vent and breast areas. Use 1 pound per 100 birds. Repeat dusting in 4 weeks, if necessary.

CHICKEN DUST BATH BOXES

To Control Northern Fowl Mites, Chicken Mites, Lice, and Fleas: Use 2½ pounds of Poultry Dusting Powder in a shallow box for 50 birds. This is a supplement to litter treatment for chicken mite and flea control. On ranges, 2 natural wallows may be substituted for each dust bath box.

Do not dust birds (or litter) within 7 days of slaughter.

CAUTION: **HARMFUL IF INHALED OR SWALLOWED** Do not breathe dust. Do not take internally. Avoid breathing of dust. *Skin contact may be harmful; avoid contact.* Wash hands before eating. Take shower or bath after work. *Wear regular long sleeved work clothing.* Change to clean clothing daily. Avoid contamination of feed, food containers and water.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Note for physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.


723B D-7153 Printed in U.S.A.
QUALITY CONTROLLED BY PURINA RESEARCH

NET WEIGHT
1 POUND



PURINA®

EPA REG.
NO. 602-71-AA

**POULTRY
DUSTING
POWDER**



CONTAINS
*SEVIN



ACCEPTED
Sep 13, 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBOTICIDE
FOR ECONOMIC POISON REGIS-
TRATION ACT
ED UNDER NO. 602-71

**FOR CONTROL OF POULTRY PESTS
ON CHICKENS, TURKEYS, DUCKS,
GEESE, GAME BIRDS AND PIGEONS**

Active Ingredients:
Carbaryl (1-naphthyl N-methylcarbamate)*..... 5%
Sulfur..... 10%
Inert Ingredients..... 85%
100%.

*Registered Trade Mark of Union Carbide Corporation
for the active ingredient carbaryl.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS**



Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Square • St. Louis, Mo. 63188