

CONTAINS
CYTHON
 INSECTICIDE
 THE PREMIUM GRADE MALATHION
 REG. T.M. - AMERICAN CYANAMID COMPANY

ACCEPTED
 UNDER THE FEDERAL
 FUNGICIDE AND
 RODENTICIDE ACT
 FOR USE ON PIGS
 ED UNDER NO. 602-87-
with comments

PURINA[®]

E P A REG. No.
 602-87-AA

MALATHION DUST

Controls: Lice and Ticks on Beef Cattle and Dry Cows • Lice on Hogs • Lice and Mites on Poultry and Pigeons • Hornflies on Dairy or Beef Cattle • Lice, Ticks and Keds on Sheep and Goats • Fleas on Dogs and Cats

ACTIVE INGREDIENT:

*Malathion

4.0%

INERT INGREDIENTS

96.0%

100.0%

*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.


**CAUTION: KEEP OUT OF REACH OF CHILDREN
 SEE OTHER CAUTIONS BELOW**

TAKE TIME



OBSERVE LABEL
 DIRECTIONS

D 5787



FOR LICE AND MITES ON POULTRY OR PIGEONS

Litter Treatment: Apply Malathion Dust to the litter at the rate of 1 pound to 50 square feet by means of a hand duster. Also dust nests, roosts, and adjacent areas. Provide clean nesting material after dust-

FOR LICE CONTROL ON HOGS; LICE AND TICKS ON BEEF CATTLE AND DRY DAIRY CATTLE:

Apply sufficient Malathion Dust by means of a hand or power duster to treat animals thoroughly. Pig pens should also be

FOR LICE AND MITES ON POULTRY OR PIGEONS

Litter Treatment: Apply Malathion Dust to the litter at the rate of 1 pound to 50 square feet by means of a hand duster. Also dust nests, roosts, and adjacent areas. Provide clean nesting material after dusting. Repeat as necessary.

Individual Treatment: Shake Malathion Dust into the feathers particularly around the vent, neck, and breast regions. Ruffle the feathers so that the powder can reach the skin. Repeat in 4 to 8 weeks or when necessary.

Caged Poultry: Use Malathion Dust in a hand plunger-type duster. Hold nozzle touching the feathers and apply one puff to both the vent and breast area.

Dust Baths: Place 1 pound of Malathion Dust in a shallow box per 50 birds. Allow hens to dust themselves freely. Repeat treatment as necessary.

FOR CHIGGERS ON POULTRY RANGES: Apply 25 pounds of Malathion Dust per acre. Make full coverage application one day prior to placing poultry on range. Repeat applications should be made every 2-3 weeks.

FOR HORN FLY CONTROL ON DAIRY OR BEEF CATTLE: Sprinkle Malathion Dust over the back and neck region at the rate of 2 ounces (4 tablespoonfuls) per animal. Repeat application at 10 to 14 day intervals, if necessary. Do not apply to dairy cattle during or less than 5 hours before milking.

FOR LICE, TICKS, AND KEDS ON SHEEP AND GOATS: Apply sufficient Malathion Dust by means of a hand or power duster to treat animals thoroughly. Repeat application after 2 or 3 weeks if needed. Do not apply to milk goats. Do not treat animals under 1 month of age.

FOR LICE CONTROL ON HOGS; LICE AND TICKS ON BEEF CATTLE AND DRY DAIRY CATTLE:

Apply sufficient Malathion Dust by means of a hand or power duster to treat animals thoroughly. Pig pens should also be thoroughly dusted. Repeat application at two week intervals if needed. Do not treat calves under one month of age. Do not apply to lactating dairy animals or non-lactating dairy animals within two weeks of freshening.

FOR FLEA CONTROL ON DOGS AND CATS

Rub Malathion Dust well into the hair coat. Do not repeat application for 7 days and only then if infestation is present. For control of fleas in dog and cat kennels, in pens, yards, lawns, and under houses apply Malathion Dust at the rate of 2 pounds per 1000 square feet using a hand duster. Remove manure and debris before dusting.

Caution: Avoid breathing of dust. Do not leave dust on skin or in eyes. Wash thoroughly. Avoid contamination of feed, food containers and water troughs. Gather eggs before dusting. Keep out of reach of children.

This product is toxic to fish and birds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Note—This container was packed full but due to shipping the contents may settle.

QUALITY CONTROLLED BY PURINA RESEARCH

NET WEIGHT 50 LB.

Manufactured by

RALSTON PURINA COMPANY

GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63188