

602-101 PM-15

10/26/81

103

Reason For Issue: Moving "Xylene" into the Inert Ingredient listing. Send final labels to EPA.

Date Issued: 5/16/89 (Yellow)

Prev. Date Issued: 4/16/87

(9-Square Logo)

FRONT PANEL

PURINA[®]
BRAND

Contains
" CIODRIN"*
Insecticide

DAIRY SPRAY
CONCENTRATE

EPA Reg. No. 602-101

(3-Row Checkerband[®])

1 PINT MAKES UP TO 10 GALLONS OF SPAY
FOR CONTROL OF: FACE FLIES--STABLE FLIES--HORN FLIES--

HOUSE FLIES--LICE--LONE STAR TICKS--WINTER TICKS

ACTIVE INGREDIENTS:

Dimethyl phosphate of alpha-methylbenzyl

3-hydroxy-cis-crotonate..... 21.5%

INERT INGREDIENTS..... 78.5%

100.0%

*Reg. T. M. Shell Chemical Company

KEEP OUT OF REACH OF CHILDREN

WARNING:

SEE OTHER WARNINGS ON BACK PANEL

PROTECT FROM FREEZING


NET 1 GAL

D-2534

NET 1 GAL
PURINA[®] DAIRY
BRAND
SPRAY CONCENTRATE

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812
St. Louis, Missouri 63166-6812 U.S.A.

Not Reviewed. Registrant claims to be in accordance with
Reregistration of Pesticide Product Guidance for Phase 2
Response, Page 2.12, Active to Inert Change in Status.



Back Panel

HOW TO USE PURINA[®] DAIRY SPRAY CONCENTRATE
BRAND

For Dairy Cattle--Beef Cattle--Sheep--Goats--Hogs

For Control of Stable Flies, House Flies, Horn Flies, and Face Flies on Dairy and Beef Cattle :

Using low pressure equipment (compression sprayer, knapsack sprayer)--Mix 1 pint of Dairy Spray Concentrate in 2-1/2 gallons of water (6-1/2 oz. per gallon). Spray to thoroughly cover all parts of the animal, including the legs, using 1 to 2 pints of spray per large animal and proportionately less for small animals.

Using high pressure equipment (hydraulic sprayer for pen or corral operations)-- Mix 1 pint of Dairy Spray Concentrate in 5 gallons of water. Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1 to 2 quarts of spray per large animal and proportionately less for smaller animals.

Repeat above application as necessary, to maintain control, but not more often than once every 7 days for either low or high pressure equipment.

To improve face fly control, add 1/2 pound of sugar per 5 gallons of spray.

Using hand atomizer sprayer --Mix 3 oz. (6 tablespoonfuls) of Dairy Spray Concentrate per quart of water. Spray to thoroughly cover all parts of the animal, including the legs. Use 1-2 fluid ounces of spray per animal daily. To improve face fly control, add 2 level tablespoons of sugar per quart of spray.

For Control of Lice, Lone Star Ticks, and Winter Ticks on Dairy Cattle Beef Cattle, Sheep, Goats and Hogs : Mix 1 pint of Dairy Spray Concentrate in 10 gallons of water. Spray animals thoroughly using up to 1 gallon of spray per large animal and proportionately less for smaller animals. Apply a second application 14 days later. Repeat applications as necessary to maintain control.

For Cattle Rubbing Devices : For the control of horn flies and face flies : Mix 1 pint of Dairy Spray Concentrate in 3 gallons of #2 fuel oil or diesel fuel oil. Install rubbing devices in areas where the animals loaf, feed, or water.

WARNING: HAZARDOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.
Hazardous or fatal if swallowed. If swallowed, induce vomiting. Call a physician immediately. Vapor harmful. Hazardous if absorbed through the skin. Atropine is antidotal.

Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Keep away from heat and open flame.

Do not apply to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphate. Do not contaminate feed, foodstuffs, or drinking water. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type approved by the U. S. Bureau of Mines for protection against Clodrin[®] insecticide. Use this material only for recommended purposes and at recommended dosages.

This product is toxic to fish and wildlife. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.

QUALITY CONTROLLED BY PURINA RESEARCH

EPA Est. No. 602-MO-1

8905 C

Printed in U.S.A.

