

025,177
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND MITICIDE ACT
 FOR ECONOMIC PEST CONTROL
 ED UNDER 602-107
 TO ATTACH TO THE LABEL

HOW TO USE CYGON 2-E SYSTEMIC INSECTICIDE

For Housefly Control in Dairy Barns, Pig Parlors, Loafing Sheds, Poultry Houses, Calf Barns, and Feedlots:

Residual Wall Spray—Mix 1 gallon of Cygon 2-E in 25 gallons of water (1 pint in 3 gallons of water). Thoroughly wet all fly resting areas to the point of run-off. One gallon of spray will cover 500 to 1000 square feet of wall or ceiling surfaces. Other areas where flies gather, such as around corrals, should be sprayed thoroughly. Repeat applications as necessary.

Remove animals and poultry from buildings when applying residual wall sprays.


Spot Sprays—Mix 1 pint of Cygon 2-E in 2½ gallons of water. Apply with compressed air or knapsack type sprayer to areas around doorways, windows, alleyways, feedrooms, and other fly resting spots. Repeat as necessary.

Good Sanitation is a necessary part of any effective fly control program.

Maggot Sprays—Mix 1 pint of Cygon 2-E in 2½ gallons of water. Apply as a coarse spray or with sprinkling can to fly breeding areas, such as poultry droppings in caged layer houses or manure piles. Repeat application as additional manure is added.

*Trademark of American Cyanamid Company
 Wayne, N. J., U.S. Patent 2,494,283

PURINA CYGON* 2-E SYSTEMIC INSECTICIDE



**Controls Houseflies in
 Dairy Barns—Pig Parlors—Loafing Sheds
 Poultry Houses—Calf Barns—Feedlots**

Active Ingredients:

Dimethoate (0,0-dimethyl S-(N-Methylcarbamoylmethyl) phosphorodithioate	23.4%
Inert Ingredients	76.6%
	100.0%

(Contains dimethoate 2 lb. per gallon)

NOT FOR USE ON ANIMALS OR POULTRY

WARNING: KEEP OUT OF REACH OF CHILDREN

SEE OTHER WARNINGS ON BACK PANEL

Manufactured By

RALSTON PURINA COMPANY
 General Offices • Checkerboard Square • St. Louis, Mo. 63188

General Outside Use—Mix 1 pint of Cygon 2-E in 3 gallons of water. To control houseflies, thoroughly spray exposed surfaces such as walls, fences, garbage and refuse containers around homes, recreation areas, animal quarters, food processing plants, warehouses, loading docks and refuse areas. Repeat as necessary.

Do not contaminate feed and foodstuffs, drinking fountains, litter and feed troughs.

Do not use in milk-processing rooms, milk houses or milk storage rooms.

Do not spray freshly whitewashed surfaces. Do not use in the home.

WARNING: VAPOR HARMFUL. HARMFUL OR FATAL IF SWALLOWED. IF SWALLOWED, INDUCE VOMITING. CALL PHYSICIAN IMMEDIATELY. CONCENTRATED MATERIAL CAUSES EYE IRRITATION.

Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep away from heat, sparks and open flame. Change clothes after spraying. Wash clothing before re-use. If symptoms of illness occur during or after spraying, see your physician. Do not spray in food processing plants. In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes; get medical attention.

Note to Physicians: This product may cause cholinesterase inhibition. Atropine is antidote.

Store Above 20° Fahrenheit

QUALITY CONTROLLED BY PURINA RESEARCH

ACCEPTED
 May 23, 1972
 U.S. DEPARTMENT OF AGRICULTURE
 FEDERAL BUREAU OF INVESTIGATION
 602-47
 TO ATTORNEY

HOW TO USE CYGON 2-E SYSTEMIC INSECTICIDE

For Housefly Control in Dairy Barns, Pig Parlors, Poultry Sheds, Slaughter Houses, Calf Barns, and Feedlots.

Residual Wall Spray—Mix 1 gallon of Cygon 2-E in 25 gallons of water (1 pint in 3 gallons of water). Thoroughly wet all fly resting areas to the point of run-off. One gallon of spray will cover 500 to 1000 square feet of wall or ceiling surfaces. Other areas where flies gather, such as around corrals, should be sprayed thoroughly. Repeat applications as necessary.

Remove animals and poultry from buildings when applying residual wall sprays.

Spot Sprays—Mix 1 pint of Cygon 2-E in 2 1/2 gallons of water. Apply with compressed air or knapsack type sprayer to areas around doorways, windows, alleys, feedrooms, and other fly resting spots. Repeat as necessary.

Good Sanitation is a necessary part of any effective fly control program.

Magnet Sprays—Mix 1 pint of Cygon 2-E in 2 1/2 gallons of water. Apply as a coarse spray or with sprinkling can to fly breeding areas, such as poultry droppings in caged layer houses or manure piles. Repeat application as additional manure is added.

Trademark of American Cyanamid Company
 Wayne, N. J., U.S. Patent 2,494,283

723A 0-2178 Printed in U.S.A.

**PURINA
 CYGON* 2-E
 SYSTEMIC
 INSECTICIDE**



Controls Houseflies in
 Dairy Barns—Pig Parlors—Lodges—Sheds
 Poultry Houses—Calf Barns—Feedlots

Active Ingredients:

Dimethoate (95-Dimethyl)	23.4%
S-(N-Methylcarbamoylmethyl) phosphorodithioate	76.6%
Inert Ingredients	100.0%

(Contains dimethoate 2 lb. per gallon)

NOT FOR USE ON ANIMALS OR POULTRY

**WARNING: KEEP OUT OF REACH OF CHILDREN
 SEE OTHER WARNINGS ON BACK PANEL**

Manufactured by
RALSTON PURINA COMPANY
 Ralston, Mo. • Chesterboard Square • St. Louis, Mo. 63177

General Outside Use—Mix 1 pint of Cygon 2-E in 3 gallons of water. To control houseflies, thoroughly spray exposed surfaces such as walls, fences, garbage and refuse containers around homes, recreation areas, animal quarters, food processing plants, warehouses, loading docks and refuse areas. Repeat as necessary.

Do not contaminate feed and foodstuffs, drinking fountains, litter and feed troughs.

Do not use in milk-processing rooms, milk houses or milk storage rooms.

Do not spray freshly whitewashed surfaces. Do not use in the home.

WARNING: VAPOR HARMFUL. HARMFUL OR FATAL IF SWALLOWED. IF SWALLOWED, INDUCE VOMITING. CALL PHYSICIAN IMMEDIATELY. CONCENTRATED MATERIAL CAUSES EYE IRRITATION.

Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep away from heat, sparks and open flame. Change clothes after spraying. Wash clothing before re-use. If symptoms of illness occur during or after spraying, see your physician. Do not spray in food processing plants. In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes. Get medical attention.

Kept in 16 oz. plastic bottle. This product may cause cholinesterase inhibition. Atropine is antidote.

Store Above 20° Fahrenheit

QUALITY CONTROLLED BY PURINA RESEARCH