

CONTENTS
1 U.S. GALLON

HOW TO USE PURINA STOCK SPRAY SPECIAL

MIXING DIRECTIONS:

Beef Cattle and Hogs: For the control of Horn Flies, Lone Star Ticks, Winter Ticks, Gulf Coast Ticks, Spinose Ear Ticks, Lice and Sarcoptic Mange Mites—Mix 1 gallon of Stock Spray Special in 100 gallons of water (2½ tablespoonfuls to 1 gallon of water) for either spraying or dipping.

Sheep and Goats: For the control of Sheep Ticks (Keds) and Lice—Mix 1 quart of Stock Spray Special in 125 gallons of water for spray solution or in 250 gallons of water for dipping vat solution.

USE DIRECTIONS:

Mix spray or dipping vat solutions thoroughly before using and keep well mixed while spraying or dipping animals. Refill dipping vats at indicated rates when fresh water must be added. Clean out entire vat when solution becomes filthy. Repeat treatment at monthly intervals, if necessary. Animals should not be treated within 30 days of slaughter. Do not spray animals confined in closed buildings. Do not spray horses or mules.

Cattle: Apply up to 1 gallon of spray solution per animal, except calves less than 6 months old. Do not directly spray or dip calves less than 6 months old. For Gulf Coast Ticks and Spinose Ear Ticks, apply

at low pressure to each ear and to head, neck and shoulder region.

Sheep and Goats: Apply 2 to 4 quarts of spray solution per animal at low pressure to lambs and adults after spring shearing. Use high pressure to penetrate fleece during fall treatment. Hold animals in dipping vats for ½ minute and duck head twice for an instant while swimming through vat.

Hogs: Spray or dip entire animal so as to thoroughly wet the skin, including the underside. Do not treat sows within 30 days of farrowing.

WARNING: Do not spray on dairy animals or in dairy barns. Do not treat lambs, kids or pigs that are in the suckling stage. Do not treat weak or emaciated animals. Do not spray horses or mules. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. May be absorbed through skin. Wash face and hands thoroughly. Avoid contamination of feed, foodstuffs, feeders, waterers, and food equipment. To protect fish and wildlife, do not contaminate streams or ponds with spray or dipping vat solution.

DANGER: Harmful or fatal if swallowed. If swallowed induce vomiting. Call physician immediately.

64128

D-9005

Printed in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH

P
STO

FOR SP
KILLS...HORN FLIES, T
Contains 25%
Active Ingredients:
*Toxaphene
**Methoxychlor, Technical
Aromatic Petroleum Deri
Inert Ingredients

*Technical chlorinated cam
**Equivalent to 19.36%
isomers and related comp

CAUTION:
SEE (

R A L S T O
GENERAL OFFICES

9-9-70
602-112

CK SPRAY SPECIAL

pressure to each ear and to head, neck and
er region.

and Goats: Apply 2 to 4 quarts of spray
n per animal at low pressure to lambs and
after spring shearing. Use high pressure to
ate fleece during fall treatment. Hold animals
ing vats for 1/2 minute and duck head twice
instant while swimming through vat.

Spray or dip entire animal so as to thoroughly
e skin, including the underside. Do not treat
within 30 days of farrowing.

ING: Do not spray on dairy animals or in
barns. Do not treat lambs, kids or pigs that
the suckling stage. Do not treat weak or
ted animals. Do not spray horses or mules.
t breathe spray mist. Do not get in eyes, on
on clothing. May be absorbed through skin.
face and hands thoroughly. Avoid contami-
of feed, foodstuffs, feeders, waterers, and
quipment. To protect fish and wildlife, do not
nitate streams or ponds with spray or dip-
at solution.

ER: Harmful or fatal if swallowed. If
owed induce vomiting. Call physician
iately.


Printed in U.S.A.

PURINA RESEARCH

CONTENTS
1 U.S. GALLON

U.S.D.A. REG. NO. 602-112

PURINA® STOCK SPRAY SPECIAL



FOR SPRAYING OR DIPPING VATS
KILLS... HORN FLIES, TICKS, LICE, FLEAS AND SARCOPTIC MANGE MITES

Contains 25% Toxaphene and 25% Methoxychlor (by volume)

Active Ingredients:	
*Toxaphene	22%
**Methoxychlor, Technical	22%
Aromatic Petroleum Derivative Solvents	48%
Inert Ingredients	8%
	<hr/> 100%

*Technical chlorinated camphene (chlorine content 67-69%).
**Equivalent to 19.36% 2,2-bis (p-methoxyphenyl) -1,1,1-trichloroethane and 2.64% other
isomers and related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS ON BACK PANEL

MANUFACTURED BY
RALSTON PURINA COMPANY
GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63199