

INSECTICIDE MIST SPECIAL

FOR FOOD PROCESSING PLANTS, MILLS, WAREHOUSES, BOX
CARS, GRANARIES, DAIRIES, AND STORAGE AREAS.




Aids in Controlling Flies, Small Flying Moths, and Roaches, Ants, Confused Flour Beetle, Cadelles, Cheese Mites, Sawtooth Grain Beetle, Red Flour Beetle, Rusty Grain Beetle.

Active Ingredients:

Pyrethrins	0.50%
Piperonyl Butoxide, Technical*	0.75%
Petroleum Oils	98.75%

*Consists of 0.60% of (butylcarbityl) (6-propylpiperonyl) ether and 0.15% of related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS BELOW



DIRECTIONS

Use INSECTICIDE MIST SPECIAL when food processing plants or mills are not in operation. Close doors and windows, turn off ventilating systems, and extinguish open flames. In poultry, meat, rabbit, egg, fish, and other food processing plants-- remove exposed products during spraying. Clean premises thoroughly by washing before operations are resumed. Repeat spraying as necessary. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

To kill exposed stages of flying insects (flies, gnats, wasps, mosquitoes, and small flying moths)-- Direct the spray mist towards the upper portions of the enclosure. Use INSECTICIDE MIST SPECIAL at the rate of 2 ounces for each 1,000 cubic feet of space. Keep doors and windows closed for 10 minutes following spraying.

To kill exposed stages of roaches, ants, confused flour beetle, sawtooth grain beetle, red flour beetle, rusty grain beetle, cadelles, and cheese mites-- Apply as a coarse wet spray into all places commonly inhabited by these insects. Contact as many insects as possible with spray. Use 1 gal-

lon of INSECTICIDE MIST SPECIAL for each 1,000 square feet of surface. Spray floors, walls, and other surfaces of bins, storage and handling areas thoroughly.

CAUTION: Harmful or fatal if swallowed. If swallowed, do not induce vomiting. Call doctor immediately. Avoid breathing of spray mist. Do not remain in treated areas and ventilate the area after treatment is completed.

Do not store near open flame. Extinguish pilot lights and open flames before space spraying. Do not apply to uncovered human foodstuffs or food products. Do not allow spray to remain on any plastic materials. Avoid contamination of milk and milk equipment.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Remove pets, birds and cover fish aquariums before spraying.

Do not re-use empty container. Destroy when empty.

QUALITY CONTROLLED BY PURINA RESEARCH

Manufactured by

RALSTON PURINA COMPANY

General Offices • Checkerboard Sq • St. Louis, Mo. 63188

EPA Est. 602-MO-1


NET 55 GALLONS

P-4536

PM 11 602-126

Purina

INSECTICIDE MIST SPECIAL



FOR FOOD PROCESSING PLANTS, MILLS, WAREHOUSES, BOX
CARS, GRANARIES, DAIRIES, AND STORAGE AREAS.


Aids in Controlling Flies, Small Flying Moths, and Roaches, Ants, Confused Flour Beetle, Caddis, Cheese Mites, Sawtooth Grain Beetle, Red
Flour Beetle, Rusty Grain Beetle.

Active Ingredients:

Pyrethrins 0.50%
Piperonyl Butoxide, Technical* 0.75%
Petroleum Oils 98.75%

*Consists of 0.60% of (butylcarbityl) (6-propylpiperonyl)
ether and 0.15% of related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS BELOW



DIRECTIONS

Use INSECTICIDE MIST SPECIAL when food processing plants or mills are not in operation. Close doors and windows, turn off ventilating systems, and extinguish open flames. In poultry, meat, rabbit, egg, fish, and other food processing plants—remove exposed products during spraying. Clean premises thoroughly by washing before operations are resumed. Repeat spraying as necessary. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

To kill exposed stages of flying insects (flies, gnats, wasps, mosquitoes, and small flying moths)—Direct the spray mist towards the upper portions of the enclosure. Use INSECTICIDE MIST SPECIAL at the rate of 2 ounces for each 1,000 cubic feet of space. Keep doors and windows closed for 10 minutes following spraying.

To kill exposed stages of roaches, ants, confused flour beetle, sawtooth grain beetle, red flour beetle, rusty grain beetle, caddis, and cheese mites—Apply as a coarse wet spray into all places commonly inhabited by these insects. Contact as many insects as possible with spray. Use 1 gal-

lon of INSECTICIDE MIST SPECIAL for each 1,000 square feet of surface. Spray floors, walls, and other surfaces of bins, storage and handling areas thoroughly.

CAUTION: Harmful or fatal if swallowed. If swallowed, do not induce vomiting. Call doctor immediately. Avoid breathing of spray mist. Do not remain in treated areas and ventilate the area after treatment is completed.

Do not store near open flame. Extinguish pilot lights and open flames before space spraying. Do not apply to uncovered human foodstuffs or food products. Do not allow spray to remain on any plastic materials. Avoid contamination of milk and milk equipment.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Remove pets, birds and cover fish aquariums before spraying.

Do not re-use empty container. Destroy when empty.

QUALITY CONTROLLED BY PURINA RESEARCH

Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63108

EPA Est. 602—MO-1

NET 55 GALLONS

P-4536 1661