

Reason For Issue: Moving "Petroleum Oils" from the Active Ingredient listing to the Inert listing. 1-25-90 '2

Date Released: 1/17/90 (Green)

Prev. Date Released: 5/15/89

(9-Square Logo)

(3-Row Checkerband)

EPA Reg. No. 602-128

Front Panel

PURINA  
BRAND

RUB-ON  
HORSE INSECTICIDE

KILLS AND REPELS

Horse Flies, Stable Flies, Horn Flies, House Flies, Deer Flies, Mosquitoes, and Gnats

AIDS IN GROOMING

(Horse Illustration)

ACTIVE INGREDIENTS:

Pyrethrins .....	0.10%
*Piperonyl butoxide, technical .....	0.20%
**Di-n-propyl isocinchomeronate .....	0.70%
***N-octyl bicycloheptene dicarboximide .....	0.35%
INERT INGREDIENTS .....	98.65%
	<u>100.00%</u>

\*Consists of 0.16% of (Butylcarbityl)(6-Propylpiperonyl) Ether and 0.04% of Related Compounds

\*\*MGK Repellent 326 U.S. Pat. No. 2,757,120

\*\*\*MGK Insecticide Synergist U.S. Pat. No. 2,576,512

Inert ingredients include Aliphatic petroleum hydrocarbons

KEEP OUT OF REACH OF CHILDREN


CAUTION

SEE OTHER CAUTIONS ON BACK PANEL

NET 1 GAL

R-8654

Manufactured By  
PURINA MILLS, INC.  
P. O. Box 66812  
St. Louis, Missouri 63166-6812



Not Revisited. Registrant must be in accordance with  
EPA's Phase 2 Guidance for Phase 2  
Response. Page 2 12. Active to Inert Change in Status.

## Back Panel

PURINA Rub-On Horse Insecticide contains a combination of insecticides and fly repellents in a blend of fine grooming oils. When applied as directed, horse flies, stable flies, horn flies, house flies, deer flies, mosquitoes, or gnats are repelled or killed. Flies not repelled stop biting and soon drop to ground and die.

Use same directions for use of this product on beef or dairy show cattle.

### DIRECTIONS FOR USE

Apply with a cloth which is slightly moistened with the product. Use 1 to 2 ounces of PURINA Rub-On Horse Insecticide per animal each day for a 3-day period. Repeat after 2 days have elapsed and every second day thereafter or when insects again disturb the animal. Briskly rub cloth against growth of hair on legs, shanks, belly, shoulders, neck and facial areas.

**COLTS and CALVES:** Treat very lightly once a day instead of attempting to build up a residual repellency by treating more heavily.

For best fly protection, apply during morning hours; for mosquitoes apply early in evening. Dusty animals should be brushed before applying PURINA Rub-On Horse Insecticide.

Keep cloth in closed metal container between treatment periods.

**CAUTION:** Do not store near an open flame. Do not wet the skin of animals. Avoid contamination of feed, foodstuffs, milk and milking utensils. Wash hands after applying.

**DANGER:** Harmful or fatal if swallowed. If swallowed do not induce vomiting. Call physician immediately.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Do not store near heat or open flame.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

EPA Est. No. 602-MO-1

QUALITY CONTROLLED BY PURINA RESEARCH

9001 L

Printed in U.S.A.

