PURINA® INSECTICIDE



Active Ingredients: Technical Chlordane*		. 25.0%
Active Ingredients: Technical Chlordane*	•••	. 25.0% . 39.5%
Inert Ingredients	• • •	10.5%
	,	100.0%

*Equivalent to 15% Octochloro-4,7-methanotetrahydroindane and 10% insecticidally active related compounds.

Not for use on animals or poultry **CAUTION:** KEEP OUT OF REACH OF CHILDREN

> See Other Warnings on Back Panel Manufactured By

RALSTON PURINA COMPANY Gen. Offices • Checkerboard Sq. • St. Louis, Mo.

HOW TO USE PURINA CHLORDANE 2-E INSECTICIDE

Crawling Pests in Farm Buildings and Household:

Mixing Directions: Mix 4 ounces (8 tablespoonfuls) of Chlordane 2-E Insecticide in 1 gallon of water.

Use Directions: Repeat applications as often as necessary to maintain effective control.

Ants, Cockroaches, Silverfish, Centipedes, Crickets, and Scorpions: Paint or spray baseboards, cracks, door and window sills, cabinets, closets, around pipe openings, and other hiding places.

Spiders: Spray webs and corners where spiders gather.

Brown Dog Ticks: Spray or paint baseboards, door and window sills, edges of carpet, behind pictures and draperies, and all other possible hiding places. Treat dogs with a Purina product labeled for tick control. DO NOT TREAT DOGS WITH THIS PRODUCT.

WARNINGS: Use on limited areas only. Do not treat entire walls or floor. Avoid spotting wallpaper and fabrics. Do not contaminate food, water, dishes, or utensils. Allow treated food and dish shelves to dry; cover with shelf paper before replacing articles. Do not allow children, pets, or animals near treated surfaces until spray has dried. Do not use in dairy barns or poultry houses.

Pests in Outdoor Areas:

Mixing Directions: Mix 4 ounces (8 tablespoonfuls) of Chlordane 2-E Insecticide in 1 gallon of water.

House Flies, Face Flies, and Stable Flies Outside of Barns: Thoroughly spray exterior surfaces of buildings and surrounding vegetation where flies rest. Apply to manure and refuse to control breeding. Use 1 gallon of mixed solution for 1000 sq. ft.

Wasps: Spray nests, particula / nest openings; around windows and areas frequented by wasps.

676D

D-1252

Boxelder Bugs: Spray ground or side of buildings where in-

WARNINGS: Use only outside for these pests. Do not use on poultry or inside poultry houses or dairy barns. Do not use on crops or grazing area.

Pests of Lawn and Turf Areas (Not Grazed):

Ants, Chinch Bugs, Japanese Beetles, White Grubs, and Chafer Beetles; Apply I pint of Chlordane 2-E Insecticide to each 1000 sq. ft. Use sufficient water for adequate coverage. Sprinkle after applying to wash insecticide into soil. Apply when insects are known to be present. A second application may be needed in 7-10 days for chinch bug control.

Sod Webworms, Mole Crickets, Cutworms, Wireworms, Armyworms, Wild Bees: Apply 1/2 pint of Chlordane 2-E Insecticide to each 1000 sq. ft. Use sufficient water for adequate coverage. Apply when insects are known to be present. Repeat if necessary.

WARNINGS: Do not treat lawns or other turf areas when people or animals are on them. Do not allow sprays to drift. to areas where they might injure people or animals or contaminate food or feed. Keep children and pets off treated areas until the insecticide has been washed off by sprinkling and the grass has completely dried. Birds feeding on treated areas may be harmed.

CAUTION: Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust and spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTIDOTE:

External—In case of spillage on skin, wash promptly with soap and water.

Internal Induce vomiting using 1 tablespoonful mustard in glass of water. Call a physician.

This product is toxic to fish and wildlife. Apply this product only as specified on this label. Rinse equipment, break empty product bottles, and dispose of wastes by burying in non-Printed in U.S.A. crop lands away from water supplies.

OUALITY CONTROLLED BY PURINA RESEARCH

602 -129