



ACCEPTED  
2/25/69  
LIBRARY

69-11-10

Active Drug Ingredients: 0.1%  
 Inert Ingredients: 99.9%

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**ADDITIONAL CAUTIONS BELOW**

**PURINA STARLICIDE COMPLETE**  
 Purina Starlicide Complete is a highly effective, fast-acting, and safe bird control agent. It is a complete feed supplement containing a potent, non-toxic, and non-flammable chemical which is effective against starlings and blackbirds. Purina Starlicide Complete is a complete feed supplement containing a potent, non-toxic, and non-flammable chemical which is effective against starlings and blackbirds. Purina Starlicide Complete is a complete feed supplement containing a potent, non-toxic, and non-flammable chemical which is effective against starlings and blackbirds.

**FOR LIVESTOCK FEEDLOTS**

Place in feeding stations or scatter Purina Starlicide Complete thinly and uniformly over dry or frozen areas of empty pens, alleyways, and other areas inaccessible to livestock. Application should be made before starlings arrive for their first morning meal. One baiting may last up to three days.

The quantity of Purina Starlicide Complete required should not exceed twenty pounds per acre for large lots (over ten acres), nor exceed fifty pounds per acre for small lots (under ten acres).

**FOR POULTRY OPERATIONS**

Place in feeding stations or scatter Purina Starlicide Complete thinly and uniformly over dry or frozen areas of empty pens, alleyways, and other areas inaccessible to poultry. Application should be made before starlings arrive for their first morning meal. One baiting may last up to three days. Because Purina Starlicide Complete is toxic to chickens, turkeys, and other poultry, it should not be used in areas accessible to poultry.

The quantity of Purina Starlicide Complete required should not exceed twenty pounds per acre for large lots (over ten acres), nor exceed fifty pounds per acre for small lots (under ten acres).

**NOTE:** Do not expect to find large numbers of dead starlings and/or blackbirds in the baited area as most die within one to three days after consuming the treated pellets. Most die at their roost and along flight lines to the feeding areas. If starlings or blackbirds continue to be a problem, assess the problem and re-baiting may be necessary. Do not re-bait within three days of application as some birds consuming a second dose of Purina Starlicide Complete may require three days to be affected.

**CAUTION**

Handle with protective gloves. Wash hands and equipment used for bait distribution thoroughly with soap and water after handling. Avoid contact with eyes, skin and clothing. If swallowed, induce vomiting and call a physician.

Do not contaminate streams, lakes or ponds with this material.

Purina Starlicide Complete must not be used in any manner that may endanger desirable and protected bird species and domestic fowl. If there is a question of hazard or legality, consult your state or local game authorities before undertaking starling control with Purina Starlicide.

**Manufactured By RALSTON PURINA COMPANY**

General Offices • Checkerboard Square • St. Louis, Missouri 63199

Printed in U.S.A.

**QUALITY CONTROLLED BY PURINA RESEARCH**

684C