

DIRECTIONS FOR USE

1. For sanitizing only.

To sanitize surfaces of buildings, equipment or utensils which have previously been thoroughly cleaned, use 3 ounces of i-Q Germicide in 5 gallons of water. Spray or swab solution on surfaces to be sanitized. Allow to drain dry without rinsing.

i-Q Germicide used at the level of 3 ounces in 5 gallons of water will pass the requirements of the official AOAC Detergent-Sanitizer Test, killing 99.999% of the test organism (E. coli) in 30 seconds.

2. For Cleaning and Disinfecting Buildings and Equipment.

Clean and disinfect in one easy step with i-Q Germicide. Use 4 teaspoonfuls of i-Q Germicide per gallon of lukewarm water, or 2 ounces in 3 gallons of water. Scrub surfaces to be cleaned and disinfected with a good brush or with a power sprayer. Start at the highest point of the building or equipment being cleaned, and work down. Let solution drain dry without rinsing.

Two ounces of i-Q Germicide in 3 gallons of water will pass the disinfectant requirements of the official AOAC Use Dilution Confirmation Method. The Staphylococcus aureus and Salmonella choleraesuis test organisms are completely killed in at least twenty-nine out of thirty tubes.

For exceptionally dirty or porous surfaces, use 4 ounces of i-Q Germicide in 3 gallons of water to clean and disinfect.

i-Q Germicide cleans and disinfects in one simple step. However, gross contamination should always be removed first (by scraping, power spraying or shoveling) before using i-Q Germicide. It is impossible to disinfect surfaces with large amounts of filth present.

3. Shoe and Boot Disinfectant Pens.

To help prevent tracking disease organisms into poultry houses, farrowing houses and dairy barns, disinfect shoes or boots in a pan containing 3 ounces of i-Q Germicide per gallon of water. Change solutions daily.

4. Deodorant.

i-Q Germicide destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab or spray on a solution of 1 ounce of i-Q Germicide per gallon of water.

APPLICATIONS: i-Q Germicide kills a broad range of bacteria, fungi and viruses responsible for such common diseases as CRD and pullorum in poultry; pig scours and calf scours.

WARNING: 1. Avoid contamination of food.

2. i-Q Germicide may cause skin irritation or damage to eyes. Avoid contact with skin and eyes. May be harmful if swallowed. In case of contact flush immediately with plenty of water. For eyes, get medical attention.


QUALITY CONTROLLED BY PURINA RESEARCH

NET CONTENTS: 1 U.S. GALLON

U.S.D.A. REG. No. 602-137

PURINA® i-Q™ GERMICIDE

Disinfectant, Deodorant and Detergent-Sanitizer with residual bacteriostatic and self-sanitizing action. For cleaning, disinfecting and sanitizing animal housing and equipment; for shoe and boot disinfection, as a deodorant.



ACTIVE INGREDIENTS:

Polyethoxy polypropoxy polyethoxy-ethanol, n-alkyl (50% C ₁₂ , 30% C ₁₁ , 17% C ₁₀ , 3% C ₁₁) dimethyl 3,4-dichlorobenzyl ammonium chloride-iodine complex*.....	13.8%
Providing: 1.50% titrable iodine and 1.50% n-alkyl dimethyl 3,4-dichlorobenzyl ammonium chlorides.....	
Phosphoric Acid.....	7.5%
Isopropyl Alcohol.....	5.0%
INERT INGREDIENTS:	73.7%

TOTAL.....100.0%

*West Chemical's Tetradine brand of Tamed Iodine.
Manufactured under the following U. S. Patents: 2,679,533 • 2,759,869
2,977,315 • 3,028,299. Additional patents pending.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL WARNINGS ON BACK PANEL

MANUFACTURED EXCLUSIVELY FOR
RALSTON PURINA COMPANY
GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63199