

2

ACCEPTED
602-1312
APR 3 1980
IN THE FEDERAL INSECTICIDE
HERBICIDE AND RODENTICIDE ACT
REGISTER

PRECAUTIONARY STATEMENTS

WARNING

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Causes eye or skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Purina I-O Concentrate is a "Controlled Iodine" disinfectant containing biodegradable detergents for industry, institution, hospital, and farm use. Recommended for disinfection of floors, walls, equipment, restrooms, laboratories, refuse areas, animal rooms and cages.

How to Mix: Blend Purina I-O Concentrate with a small amount of water first; then dilute out to volume.

Official Disinfectant: Tuberculocidal-Permitted for use by the Animal Health Programs, Veterinary Service, Animal and Plant Health Service, USDA, in official disinfection programs involving domestic animal housing and transportation facilities. Add 1 ounce (2 tablespoonfuls) of Purina I-O Concentrate to each 2 gallons of water.

Poultry and Livestock Buildings, Dog Kennels, Laboratories, Zoos: Remove all animals and feeds from premises, cars, trucks, coops, and crates. Remove all litter and manure from floors, walls, and other surface areas of facilities. Empty all troughs, racks, and other feeding and watering appliances. Add 1 ounce (2 tablespoonfuls) of Purina I-O Concentrate to each 4 gallons

of water and saturate all surface areas. To disinfect porous or hard-to-clean surfaces, use product at the rate of 1 ounce in each 2 gallons of water. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers. Ventilate buildings and other closed areas, and allow the disinfectant solution to dry or be absorbed before animals are housed. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be thoroughly scrubbed with detergents and rinsed with potable water prior to re-use.

Food Contact Surfaces: To sanitize surfaces after they have been thoroughly cleaned, use 1 ounce (2 tablespoonfuls) of Purina I-O Concentrate to each 12 gallons of water. At this level, no potable water rinse is required.

For Cleaning Sows: Before placing sows in farrowing pens or crates, scrub down flanks and udder with lukewarm solution of 1 ounce (2 tablespoonfuls) Purina I-O Concentrate to each 2 gallons of water. Use a soft-bristled brush. Let sows drain dry, and do not rinse until just before farrowing.

For Shoe and Boot Sanitizing Pans: Use Purina I-O Concentrate at the rate of 1 ounce (2 tablespoonfuls) to each 2 gallons of water to help reduce tracking of disease organisms into poultry houses, farrowing houses, and other animal quarters. Change solutions daily.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Disposal - Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Container Disposal - Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

Consult Federal, State, or local disposal authorities for approved alternative procedures such as limited open burning.

DE
ICT
TEIL

1 saturate all surface areas. To disinfect porous or hard-to-clean surfaces, use a rate of 1 ounce in each 2 gallons of water. Immerse all halters, ropes, and other equipment used in handling and restraining animals as well as forks, shovels, and ventilate buildings and other closed areas, and allow the disinfectant solution to dry before animals are housed. All treated feed racks, mangers, troughs, automatic waterers, and waterers must be thoroughly scrubbed with detergents and rinsed with water prior to re-use.

Surfaces: To sanitize surfaces after they have been thoroughly cleaned, use 1 tablespoonful of Purina i-O Concentrate to each 12 gallons of water. At this level no rinsing is required.

Sows: Before placing sows in farrowing pens or crates, scrub down flanks and udders with a warm solution of 1 ounce (2 tablespoonfuls) Purina i-O Concentrate to each 2 gallons of water. Use a soft-bristled brush. Let sow drain dry, and do not rinse until just before farrowing.

Boot Sanitizing Pans: Use Purina i-O Concentrate at the rate of 1 ounce (2 tablespoonfuls) to each 2 gallons of water to help reduce tracking of disease organisms into poultry houses, and other animal quarters. Change solutions daily.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not use empty container.



Disposal—Pesticide: spray mixture or residue that cannot be used or chemically treated should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Disposal—Triple rinse (or equivalent) and dispose in an incinerator or landfill for pesticide containers, or bury in a safe place.

Follow Federal, State, or local disposal authorities for approved alternative procedures such as incineration.

10. 602-Mo-1

Printed in U.S.A.



ACCEPTED
602-138
APR 3 1960
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR SOLOSOND POISON REGISTERED
UNDER NO.

KEEP FROM CHILDREN
GARDEN EXCLUSIVE
(See label and directions for use)
KEEP FROM PETS
(If treated let them, etc. well before)

NET CONTENTS: 1 GAL




PURINA[®] I-O[®] CONCENTRATE

EPA Reg. No. 602-138

"CONTROLLED 100%
General Purpose Disinfectant

A detergent germicide for sanitizing and disinfecting environmental surfaces. Always wear eye protection, and always use best sanitizing practices.



ACTIVE INGREDIENTS:

2,4-Dichloro-1,3-bis(4-chlorophenyl)butane (containing 2,4-D) 27.0%

Sodium hypochlorite (containing 5.25% available chlorine) 6.0%

Water 66.8%

27.0%
6.0%
66.8%
100.0%

U.S. Patent Nos. 2,931,777-3,008,268

ACCEPTED
602-138
APR 8 1988
FEDERAL BUREAU OF INVESTIGATION
DEPARTMENT OF JUSTICE
WASHINGTON, D.C.

KEEP OUT OF REACH OF CHILDREN
WARNING
CAUSES SKIN AND EYE IRRITATION
(See first aid and other precautions below.)

KEEP FROM FREEZING
(If frozen, let thaw, mix well before using.)

NET CONTENTS: 1 GAL

Manufactured by
RALSTON PURINA COMPANY
General Office • Checkerboard Building • St. Louis, Missouri 63108

40