

**HOW TO USE
PURINA TOMATO AND GARDEN DUST**

This product may be used as either a Dust or Spray

Apply thoroughly to upper and lower surfaces of all foliage. Protect plants from early insect and disease damage. Repeat at 7 day intervals or as necessary. Do not apply within 14 days of harvest.

DUST

Dust on plants covering leaves evenly and lightly. Apply when air is still and plants are dry. Best coverage is obtained by use of a good dust gun.

SPRAY

Mix 4 ounces (12 tablespoonfuls) in 1 1/4 gallons of water. Keep agitated while spraying. Apply as a wetting mist with either a pressure or hand type sprayer.

ASPARAGUS: Asparagus beetle, thrip. **Disease**—rust. Apply only on asparagus ferns after spears have been harvested.

BEANS: Aphids, bean leaf beetle, Mexican bean beetle, cucumber beetle, flea beetle, Japanese beetle, leafhopper, velvetbean caterpillar, western bean cutworm, armyworm, corn earworm, Lima bean pod borer, stink bug, tarnished plant bug, lygus bug, spider mite. **Disease**—anthracnose, rust, downy mildew.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: Aphids, flea beetle, harlequin bug, armyworm, cabbage looper, imported cabbage worm. **Diseases**—downy mildew, alternaria leaf spot.

CUCUMBER, MELONS, SQUASH, PUMPKIN: Aphids, cucumber beetle, flea beetle, leafhopper, melonworm, pickleworm, squash bug, squash vine borer, spider mite. **Diseases**—downy mildew, alternaria blight, anthracnose, gummy stem blight.

TOMATO, EGGPLANT, PEPPER, PEAS: Aphids, tomato fruit worm, tomato hornworm, flea beetle, grasshopper, leafhopper, European corn borer, fall armyworm, lace bug, tarnished plant bug, stink bug, spider mite, pea weevil.

Diseases—early and late blight, anthracnose, stemphylium and septoria leaf spots, cladosporium grey mold.

LETTUCE, CARROTS, BEETS, KALE, SPINACH, TURNIPS: Aphids, flea beetle, leafhoppers, cabbage looper, imported cabbage worm, harlequin bug, armyworm, corn earworm, spittlebug, tarnished plant bug, stink bug, grasshopper, Japanese beetle. **Diseases**—downy mildew, leaf spot, cercospora leaf spot.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Note for Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal for carbaryl and malathion.

ANTIDOTE: If swallowed give emetic of salt in warm water. Wash from skin with soap and water; for eyes get medical attention.

Note: Do not feed treated foliage, fruit or pods to dairy animals or animals being finished for slaughter. Toxic to honey bees; do not apply to open blossoms.

Toxic to fish; keep out of any body of water. Use only as specified on this label.

KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS

QUALITY CONTROLLED BY PURINA RESEARCH


703E

G-9503

Printed in U.S.A.

TAKE TIME

OBSERVE LABEL
DIRECTIONS



U.S.D.A. REG. NO.
602-150

1-22-71
602-150

**PURINA
TOMATO &
GARDEN DUST**

CONTROLS INSECTS AND PLANT DISEASES



Active Ingredients:

*Malathion.....	4.00%
**Carbaryl (1-Naphthyl N-methylcarbamate)....	3.00%
Zineb (Zinc ethylene bisdithiocarbamate).....	3.90%
Inert Ingredients.....	89.10%
	<u>100.00%</u>

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

**Sevin Registered Trade-mark of Union Carbide Corporation
U.S. Pat. No. 2,903,478

CAUTION KEEP OUT OF THE REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET WT. 16 OZ. (1 LB.)

Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Square • St. Louis, Mo. 63199