

RALSTON PURINA CO. ST LOUIS MO

TAKE TIME



OBSERVE LABEL DIRECTIONS

NET WEIGHT SHOWN ON BAG U.S.D.A. # 602-152

# PURINA® CHECK-FLY™ RANGE CHOW® 1-3 MIX

Medicated

To Prevent Development of Horn Fly and Face Fly Maggots in Cattle Manure; to Retard Parasitic Egg Production and Larval Development of Common Stomach Worms (haemonchus), Nodular Worms (oesophagostomum), and Hookworms (bunostomum) in Cattle by Interrupting Life Cycle. Aids in the Prevention of Foot Rot in Cattle and Soft Tissue Lumpy Jaw; Nutritional Source of Iodine.

**Pesticidally Active Drug Ingredients:**

Phenothiazine\* . . . . . 0.31 %

**Pesticidally Inert Ingredients** . . . . . 99.682%

**Active Drug Ingredients:**

Ethylene Diamine Dihydriodide . . . . . 0.008%

\*Average particle size of phenothiazine N.F. 6-8 microns.

**CAUTION: Keep Out of Reach of Children**

Take Time - Observe Label Directions on Reverse Side

Manufactured by

**RALSTON PURINA COMPANY**

Gen. Offices, St. Louis, Mo. 63199

**GUARANTEED ANALYSIS**

Crude protein not less than . . . . . 15.0%

Includes not more than 6.0% equivalent protein from non-protein nitrogen

Crude fat not less than . . . . . 1.0%

Crude fiber not more than . . . . . 11.0%

Calcium (Ca) not less than . . . . . 1.0%

Calcium (Ca) not more than . . . . . 2.0%

Phosphorus (P) not less than . . . . . 0.4%

Iodine (I) not less than . . . . . 0.0001%

Salt (NaCl) not less than . . . . . 23.0%

Salt (NaCl) not more than . . . . . 27.0%

**INGREDIENTS**

Soybean meal, ground yellow corn, ground grain sorghums, cane molasses, alfalfa meal, BHT (a preservative), wheat middlings, urea, low fluorine rock phosphate, salt, iodized salt, calcium carbonate, iron oxide, manganous oxide, copper oxide, cobalt carbonate, zinc oxide. A-3234 ●



CHECK-FLY RANGE CHOW 1-3 MIX (Medicated) M

ACCEPTED WITH COMMENTS

**DIRECTIONS**

Feed at the rate of 1.5 pounds per head per day.  
Maintain a supply of clean water near feeding areas.

**WARNING**

DO NOT FEED TO LACTATING DAIRY ANIMALS. OCCASIONALLY, AN ANIMAL MAY BE SENSITIVE TO PHENOTHIAZINE RESULTING IN SYMPTOMS SUCH AS TEMPORARY BLINDNESS AND LOSS OF SKIN OR HAIR. IF TOXIC SYMPTOMS DEVELOPE, TAKE ANIMALS OFF DRUG AND AVOID EXPOSURE TO SUNLIGHT FOR SEVERAL DAYS. DO NOT ADMINISTER TO ANIMALS SHOWING SYMPTOMS OF ACUTE RESPIRATORY CONDITIONS.

**NOTE**

This product kills horn fly and face fly maggots in manure but does not control existing adult flies. If necessary, spray cattle for maximum protection. Because face flies migrate from farm to farm, it is essential that a larvicide program be employed on an area basis to assist in reducing the adult face fly population.

A-3234

603-509