

RALSTON PURINA CO. ST LOUIS MO

TAKE TIME



OBSERVE LABEL DIRECTIONS

NET WEIGHT SHOWN ON BAG U.S.D.A. #602-153

**PURINA® CHECK-FLY™  
RANGE CHOW® 1-2 MIX**

Medicated

To Prevent Development of  Horn Fly and  Face Fly  Maggots in Cattle Manure; to  Retard Parasite Egg Production and Larval Development of Common Stomach Worms (haemaphys), Nodular Worms (oesophagostomum), and Hookworms (bunostomum) in Cattle by Interrupting Life Cycle. Aids in the Prevention of Foot Rot in Cattle and Soft Tissue Lumpy Jaw; Nutritional Source of Iodine.

**Pesticidally Active Drug Ingredients:**

Phenothiazine\* ..... 0.31 %  
Pesticidally Inert Ingredients ..... 99.682%

**Active Drug Ingredients:**

Ethylene Diamine Dihydriodide ..... 0.008%

\*Average particle size of phenothiazine N.F. 6-8 microns.

**CAUTION:** Keep Out of Reach of Children

**Take Time - Observe Label Directions on Reverse Side**

Manufactured by

**RALSTON PURINA COMPANY**

Gen. Offices, St. Louis, Mo. 63199


**GUARANTEED ANALYSIS**

Crude protein not less than ..... 13.0%  
Includes not more than 3.9% equivalent protein from non-protein nitrogen  
Crude fat not less than ..... 1.0%  
Crude fiber not more than ..... 10.0%  
Calcium (Ca) not less than ..... 1.0%  
Calcium (Ca) not more than ..... 2.0%  
Phosphorus (P) not less than ..... 0.4%  
Iodine (I) not less than ..... 0.0001%  
Salt (NaCl) not less than ..... 31.0%  
Salt (NaCl) not more than ..... 36.0%

**INGREDIENTS**

Soybean meal, cottonseed meal, ground yellow corn, ground grain sorghums, cane molasses, grain sorghums sugar molasses, alfalfa meal, dehydrated alfalfa meal preserved with ethoxyquin, rice bran, BHT (a preservative), wheat middlings, urea, low fluorine rock phosphate, salt, iodized salt, calcium carbonate, iron oxide, manganese sulfate, manganous oxide, copper oxide, cobalt carbonate, zinc oxide.

CHECK-FLY RANGE CHOW 1-2 MIX (Medicated)



#### **DIRECTIONS**

Feed at the rate of 1.5 pounds per head per day.  
Maintain a supply of clean water near feeding areas.

#### **WARNING**

DO NOT FEED TO LACTATING DAIRY ANIMALS. OCCASIONALLY, AN ANIMAL MAY BE SENSITIVE TO PHENOTHIAZINE RESULTING IN SYMPTOMS SUCH AS TEMPORARY BLINDNESS AND LOSS OF SKIN OR HAIR. IF TOXIC SYMPTOMS DEVELOPE, TAKE ANIMALS OFF DRUG AND AVOID EXPOSURE TO SUNLIGHT FOR SEVERAL DAYS. DO NOT ADMINISTER TO ANIMALS SHOWING SYMPTOMS OF ACUTE RESPIRATORY CONDITIONS.

#### **NOTE**

This product kills horn fly and face fly maggots in manure but does not control existing adult flies. If necessary, spray cattle for maximum protection. Because face flies migrate from farm to farm, it is essential that a larvicide program be employed on an area basis to assist in reducing the adult face fly population.

**ACCEPTED**

**JUL 3 1 1969**

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 602-153 SUBJECT  
TO ATTACHED COMMENTS.

**DIRECTIONS**

Feed at the rate of 1.5 pounds per head per day  
Maintain a supply of clean water near feeding areas.

**WARNING**

DO NOT FEED TO LACTATING DAIRY ANIMALS. OCCASIONALLY, AN ANIMAL MAY BE SENSITIVE TO PHENOTHIAZINE RESULTING IN SYMPTOMS SUCH AS TEMPORARY BLINDNESS AND LOSS OF SKIN OR HAIR. IF TOXIC SYMPTOMS DEVELOPE, TAKE ANIMALS OFF DRUG AND AVOID EXPOSURE TO SUNLIGHT FOR SEVERAL DAYS. DO NOT ADMINISTER TO ANIMALS SHOWING SYMPTOMS OF ACUTE RESPIRATORY CONDITIONS.

**NOTE**

This product kills horn fly and face fly maggots in manure but does not control existing adult flies. If necessary, spray cattle for maximum protection. Because face flies migrate from farm to farm, it is essential that a larvicide program be employed on an area basis to assist in reducing the adult face fly population.

