
TAKE THME



PURINA' YARD & PATIO FOG

FOR QUICK KILL OF MOSQUITOES



HOW TO USE PURINA YARD & PATIO FOG

Directions For Outdoor Fogging Applications: Purina Yard and Patio Fog is a ready-to-use solution for use in foggers to control mosquitoes. Do not apply as an indoor space spray or fog.

Apply 8 ozs. ($\frac{1}{2}$ pint) for the average home yard. One gallon will treat an acre. Fog thoroughly mosquito breeding or resting areas such as bushes, heavy foliage and refuse storage areas.

For best results fog in the evening after sunset. Wind velocity should be under 4 miles per hour. The area to be fogged should always be downwind from fogging equipment.

RESTRICTIONS: Keep all persons out of the area while fogging and until the insecticide has dissipated. Keep out of any body of water. Apply only as specified on this label.

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with soap and warm water after handling. Wash clothing with soap and hot water before re-use.

Avoid contamination of feed, food, or food-handling equipment. If food handling equipment becomes contaminated, clean thoroughly before re-use. Keep out of reach of children.

Do not use, pour or store near heat or open flame. Do not fog in presence of open fire or flame.

TO PHYSICIAN: ANTIDOTE—Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Container Disposal: Do not re-use empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

QUALITY CONTROLLED BY PURINA RESEARCH G-9795 6910B

Printed in U.S.A.




ACC UNCEN TH PUNRIOIDE POR ECONON FT DEDEN M



FOR QUICK KILL OF MOSQUITOES





YARD & PATIO FOG

jing Applications:

is a ready-to-use solution i mosquitoes. Do not apply fog.

e average home yard. One Fog thoroughly mosquito ch as bushes, heavy foliage

evening after sunset. Wind miles per hour. The area to be downwind from fogging

prsons out of the area while licide has dissipated. Keep Apply only as specified on

owed, inhaled, or absorbed in eyes, on skin, or on clothnist. Wash thoroughly with r handling. Wash clothing ore re-use.

ed, food, or food-handling g equipment becomes cony before re-use. Keep out


ear heat or open flame. Do fire or flame.

-Administer atropine suloses. Repeat as necessary -PAM is also antidotal and junction with atropine.

t re-use empty container. crushing. Bury or discard


BY PURINA RESEARCH Printed in U.S.A.





PURINA' YARD & PATIO








CAUTION: KEEP OUT OF REACH OF CHILDREN SEE OTHER WARNINGS ON BACK PANEL

NET CONTENTS: 64 FL. OZ. (1/2 GAL.)

Manufactured by **RALSTON PURINA COMPANY** General Offices . Checkerboard Square . St. Louis, Missouri 63199







HOW TO USE PURINA YARD & PATIO FOG

Directions For Outdoor Fogging Applications:

Purina Yard and Patio Fog is a ready-to-use solution for use in foggers to control mosquitoes. Do not apply as an indoor space spray or for.

Apply 8 ozs. (1) pint) for the average home yard. One gallon will treat an acre. Fog thoroughly mosquito breeding or resting areas such as bushes, heavy foliage and refuse storage areas.

For best results fog in the evening after sunset. Wind velocity should be under 4 miles per hour. The area to be fogged should always be downwind from fogging equipment.

RESTRICTIONS: Keep all persons out of the area while fogging and until the insecticide has dissipated. Keep out of any body of water. Apply only as specified on this label.

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with soap and warm water after handling. Wash clothing with soap and hot water before reluse

Avoid contamination of feed, food, or food-handling equipment. If food handling equipment becomes contaminated, clean thoroughly trefore relase. Keep out of reach of children

Do not use, pour or store near heat or open flame. Do not fog in presence of open fire or flame.

TO PHYSICIAN: ANTIDOTE-Administer atropine subfate in large therapeutic doses. Repeat as necessary to the point of tolerance 2 PAM is also antidotal and may be administered in conjunction with atropine

Container Disposal: Do not re-use empty container Destroy it by perforating or crushing. Bury or discard in a safe place.

QUALITY CONTROLLED BY PURINA RESEARCH

6410B

G-9795

APPENDED AT A

