

HOW TO USE SEVIN* INSECTICIDE SPRAY

MIXING DIRECTIONS: Use 2 level tablespoonfuls per gallon of water. For pressure sprayers add directly to water; shake or agitate thoroughly. For garden hose sprayers make a slurry in bottle by mixing required product and water; agitate mixture frequently for uniform coverage.

USE DIRECTIONS: Spray foliage, buds and flowers when insects or their damage appear. Respray weekly or as needed.

FLOWERS, ROSES, SORBUS, TREES, EVERGREENS: Japanese beetle, rose slug, flea beetle, leaf roller, leaf tier, elm leaf beetle, mimosa webworm, tent caterpillar, June bug, bagworm, gipsy moth and certain other insects. **Note:** Do not use on Boston Ivy.

VEGETABLES: Beans and Peas—Mexican bean beetle, bean leaf beetle, Japanese beetle. **Beet, Cabbage, Cauliflower, Carrots, Kohlrabi, Turnip**—Flea beetle. **Cucumber, Squash, Pumpkin**—pickleworm, cucumber beetle, squash bug, squash vine borer. **Potato**—Colorado and three-lined potato beetles. **Tomato**—hornworm, fruitworm. For fruitworm, apply on foliage when fruit is setting; repeat in 2 and 4 weeks.

Do not apply within 3 days of harvest. Do not apply to beet or turnip tops used for food or feed within 14 days of harvest.

FRUITS: Peach and Apricot—Japanese beetle, red-banded leaf roller, Oriental fruit moth, stink bug. For Oriental fruit moth apply 3 times at 10 day intervals beginning at shuck split. For stink bugs apply at petal fall and 2 or 3 times 10 days apart. **Plum**—Forbes and San Jose scale. Apply when crawlers are active. **Cherry**—Japanese beetle, rose chafer. **Grapes**—Japanese beetle, red-banded leaf roller, grape berry moth. **Raspberry, Blackberry, Boysenberry**—Japanese beetle, rose chafer, stink bug. **Strawberry**—leaf roller.

Do not spray peach, plum, cherry or strawberry within 1 day; apricot within 3 days; and berries within 7 days of harvest.

LAWN INSECTS—Sod webworm (lawn moth), fall armyworm, chinch bug (except in Florida), fleas, ants, millepedes, earwigs.

Mow lawn; water thoroughly. Mix 1 tablespoonful per gallon of water; apply 1 gallon of mixture per 25 sq. ft. immediately after watering lawn. Repeat 2 to 3 weeks later if needed.


CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing spray mist. Skin contact may be harmful. Avoid contact. Wash hands before eating. Change to clean clothing daily.

Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ANTIDOTE: If swallowed give emetic of salt in warm water. Wash from skin with soap and water; for eyes get medical attention.

Note: Do not feed treated foliage, fruit or pods to dairy animals or animals being finished for slaughter. Toxic to honey bees; do not apply to open blossoms.

Use only as specified on this label.



1-29-70
602-162

PURINA SEVIN* INSECTICIDE SPRAY

FOR FLOWERS • VEGETABLES • FRUITS
ORNAMENTALS • LAWNS

Active Ingredient:	
Carbaryl (1-Naphthyl methylcarbamate)....	50.0%
Inert Ingredients.....	50.0%
	100.0%

*Trademark of Union Carbide Corporation for the active ingredient 1-Naphthyl methylcarbamate. U.S. Pat. No. 2,903,478

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON BACK PANEL

NET. WT. 8 OZ.

Distributed by
RALSTON PURINA COMPANY
General Offices • Checkerboard Square • St. Louis, Missouri 63199