

HOW TO USE PURINA ROACH BAIT

Purina Roach Bait is a ready-to-use product, effective as an aid in controlling German, Oriental and American species of cockroaches.

DIRECTIONS:

For Use in Homes, Apartment Buildings, Stores, Restaurants and Warehouses—Apply bait to floors near baseboards, in closets, under sinks and refrigerators and in other areas where roaches may hide. Scatter lightly so that bait is barely visible—use 1 level teaspoonful for 25 square feet or 4 ozs. per 1000 sq. ft. of floor space. Use fresh bait at 7 day intervals if needed.

For use By Pest Control Operators or Other Trained Personnel in Poultry, Meat, Rabbit and Seafood Packing Plants; Egg Products and Food Processing Plants—Apply bait on paper, pasteboard or other material that will permit removal. Bait may also be used in rat and mouse bait stations to which roaches are attracted. Locate these bait stations on floor near walls, in storage areas and other areas where roaches have been observed. Maintain strict account of location and number of bait stations. Do not apply in edible product areas of food processing plants. Use fresh bait at 7 day intervals if needed.

For Use in Animal Research Facilities—The use of Roach Bait in and around research animal quarters will not induce changes in liver microsomal enzyme activity in these animals.

RESTRICTIONS: Do not apply to animals. Do not treat portions of buildings where animals can feed on bait. Apply only in areas inaccessible to children and pets. Do not apply in the food preparation areas of restaurants.

CAUTION: May be harmful if swallowed or absorbed through the skin. Do not contaminate feed or food. Avoid skin contact. Wash thoroughly after using. Any product visible after treatment should be brushed into cracks or crevices or otherwise removed. Keep away from children and pets.


If poisoning occurs, get prompt medical aid. Atropine sulfate is antidotal.

Do not re-use empty container. Destroy container by perforating and bury or discard in a safe place.


725 E

D-8624

Printed in U.S.A.



PURINA[®] ROACH BAIT



Active Ingredients:
o-Isopropoxyphenyl methylcarbamate*..... 2.00%
Boric Acid..... 20.00%
Inert Ingredients..... 78.00%

*U.S. Patent No. 3,111,539

ACCEPTED

NOV 15 1972

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
NO. 602-164
SEE ATTACHED COMMENTS

TAKE TIME



OBSERVE LABEL
DIRECTIONS


CAUTION: KEEP OUT OF REACH OF CHILDREN
See Other Cautions On Back Panel

NET WEIGHT: 16 OZ. (1 lb.)

Manufactured By

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Missouri 63188



PURINA[®] ROACH BAIT

HOW TO USE PURINA ROACH BAIT

Purina Roach Bait is a ready-to-use product, effective as an aid in controlling German, Oriental and American species of cockroaches.

DIRECTIONS:

For Use in Homes, Apartment Buildings, Stores, Restaurants and Warehouses—Apply bait to floors near baseboards, in closets, under sinks and refrigerators and in other areas where roaches may hide. Scatter lightly so that bait is barely visible—use 1 level teaspoonful for 25 square feet or 4 ozs. per 1000 sq. ft. of floor space. Use fresh bait at 7 day intervals if needed.

For use By Pest Control Operators or Other Trained Personnel in Poultry, Meat, Rabbit and Seafood Packing Plants; Egg Products and Food Processing Plants—Apply bait on paper, pasteboard or other material that will permit removal. Bait may also be used in rat and mouse bait stations to which roaches are attracted. Locate these bait stations on floor near walls, in storage areas and other areas where roaches have been observed. Maintain strict account of location and number of bait stations. Do not apply in edible product areas of food processing plants. Use fresh bait at 7 day intervals if needed.

For Use in Animal Research Facilities—The use of Roach Bait in and around research animal quarters will not induce changes in liver microsomal enzyme activity in these animals.

RESTRICTIONS: Do not apply to animals. Do not treat portions of buildings where animals can feed on bait. Apply only in areas inaccessible to children and pets. Do not apply in the food preparation areas of restaurants.

CAUTION: May be harmful if swallowed or absorbed through the skin. Do not contaminate feed or food. Avoid skin contact. Wash thoroughly after using. Any product visible after treatment should be brushed into cracks or crevices or otherwise removed. Keep away from children and pets.

If poisoning occurs, get prompt medical aid. Atropine sulfate is antidotal.

Do not re-use empty container. Destroy container by perforating and bury or discard in a safe place.

D-8624

725 E

Printed in U.S.A.

Active Ingredients:
o-Isopropoxyphenyl methylcarbamate*..... 2.00%
Boric Acid.....20.00%
Inert Ingredients.....78.00%
*U.S. Patent No. 3,111,539

TAKE TIME



OBSERVE LABEL
DIRECTIONS

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Other Cautions On Back Panel

NET WEIGHT: 16 OZ. (1 lb.)

Manufactured By

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Missouri 63188

QUALITY CONTROLLED BY PURINA RESEARCH