

REGISTERED

INSECTICIDE
FEDERAL ACT
REGISTERED

How To Use PURINA CHLORDANE INSECTICIDE-45E

LAWN AND TURF INSECTS (Areas Not Grazed):
Chiggers, Fleas, Ticks, Sod Webworms (Lawn Moths),
Mole Crickets, Cutworms, Armyworms, Wild Bees,
Wireworms.

Use 1 ounce (2 tablespoonfuls) per gallon of water to cover 500 square feet of lawn. Apply when insects are known to be present. Repeat if necessary.

ANTS (including Fire and Harvester); Larvae of Japanese Beetle, Chafer Beetle, White-fringed Beetle, and Green Jung Beetle (White Grubs); Centipedes, Millipedes.

Use 2 ounces (4 tablespoonfuls) per gallon of water to cover 500 square feet of lawn. Sprinkle after applying to wash insecticide into soil. Apply when insects are known to be present. **For ant mounds** break surface of mound and saturate thoroughly. Repeat in 2 weeks if necessary.

WARNINGS: Do not treat lawns or other turf when people or animals are on them. Do not allow spray to drift to areas where they might injure people or animals or contaminate food or feed. Keep children and pets off treated areas until the insecticide has been washed off by sprinkling and the grass has completely dried.

CAUTION: Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTIDOTE:

External—In case of spillage on skin, wash promptly with soap and water.


Internal—Induce vomiting using 1 tablespoonful mustard in glass of water. Call a physician.

QUALITY CONTROLLED BY PURINA RESEARCH

698D

G-5223

Printed in U.S.A.



U.S.D.A. Reg. No.
602-166

CHLORDANE INSECTICIDE-45E

45% Emulsifiable Concentrate

Active Ingredients:

*Chlordane, Technical.....	45.0%
Aliphatic Petroleum Hydrocarbons.....	42.5%
Xylene.....	4.0%
Inert Ingredients.....	8.5%
	100.0%

*Equivalent to 27% Octachloro-4, 7-methanotetrahydroindane and 18% related compounds.

**TO BE MIXED WITH WATER
FOR LAWN AND HOUSEHOLD USES**

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANELS FOR ADDITIONAL CAUTIONS**

NET: 32 FL OZ. (1 QT.)

Manufactured by

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Missouri 63199

TERMITES: Dig a trench along out but not below the top of the footing. Apply 2 gallons of 5 linear feet where trench is not over 18 inches deep. For trenches over 18 inches deep apply 5 linear feet. For pre-construction mixed solution per 10 square feet of

HOUSEHOLD PESTS (FOR SPOT APPLICATION): Mix 2 ounces (4 tablespoonfuls) in water. Repeat application as often as necessary for effective control.

Ants, Cockroaches, Silverfish, Centipedes, Scorpions—spray or paint baseboards, sills, cabinets, closets, around pipe and other hiding places.

Spiders: Spray webs and corners where

Brown Dog Ticks: Spray or paint baseboards, edges of carpet, behind pictures and other possible hiding places. **DO NOT MIX WITH THIS PRODUCT.**

Wasps: Spray nests, particularly near windows and areas frequented by wasps.


Boxelder Bugs: Spray around or side of insects crawl. Repeat application as necessary.

WARNINGS: Use on limited areas of walls or floor. Avoid spots on fabrics. Do not contaminate food or utensils. Allow treated dish and food to cover with shelf paper before replacing. Do not allow children, pets or animals near until spray has dried. Do not use on poultry houses.

This product is toxic to fish and many other forms of life. Apply this product only to the areas indicated on this label.

REGISTERED

U.S. DEPT. OF AGRICULTURE
REGISTERED



U.S.D.A. Reg. No.
602-166

CHLORDANE INSECTICIDE-45E

45% Emulsifiable Concentrate

Active Ingredients:	
*Chlordane, Technical.....	45.0%
Aliphatic Petroleum Hydrocarbons.....	42.5%
Xylene.....	4.0%
Inert Ingredients.....	8.5%
	100.0%

*Equivalent to 27% Octachloro-4, 7-methanotetrahydroindane and 18% related compounds.

**TO BE MIXED WITH WATER
FOR LAWN AND HOUSEHOLD USES**

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANELS FOR ADDITIONAL CAUTIONS**

NET: 32 FL OZ. (1 QT.)

Manufactured by

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Missouri 63199

Use PURINA INSECTICIDE-45E

SPOTS (Areas Not Grazed):

Sod Webworms (Lawn Moths),
Grasshoppers, Armyworms, Wild Bees,

1/2 teaspoonfuls) per gallon of water to
be applied to lawn. Apply when insects are
active. Repeat if necessary.

(Harvester); Larvae of Japanese
White-fringed Beetle, and Green
Grasshoppers; Centipedes, Millipedes.

1/2 teaspoonfuls) per gallon of water to
be applied to lawn. Sprinkle after applying
to soil. Apply when insects are
active. For ant mounds break surface of
mound thoroughly. Repeat in 2 weeks if

on lawns or other turf when people
are present. Do not allow spray to drift to
injure people or animals or con-
taminant. Keep children and pets off
of treated area. Insecticide has been washed off
of grass has completely dried.

Not allowed. Contact with skin can
cause irritation. Avoid breathing spray mist. In
case of contact with skin, wash with soap and water.
Do not eat, drink, or use tobacco, feed and foodstuffs.

In case of irritation on skin, wash promptly

For more information, contact your dealer or
write using 1 tablespoonful must be
consulted by a physician.

PRODUCED BY PURINA RESEARCH

223

Printed in U.S.A.

TERMITES: Dig a trench along outside of foundation
but not below the top of the footing. Mix 1 quart in 12
gallons of water. Apply 2 gallons of mixed solution per
5 linear feet where trench is not over 18 inches deep.
For trenches over 18 inches deep apply 4 gallons per
5 linear feet. For pre-construction, apply 1 gallon of
mixed solution per 10 square feet of ground.

HOUSEHOLD PESTS (FOR SPOT APPLICATION ONLY):
Mix 2 ounces (4 tablespoonfuls) in 1 quart of water.
Repeat application as often as necessary to maintain
effective control.

**Ants, Cockroaches, Silverfish, Centipedes, Crickets,
Scorpions**—spray or paint baseboards, cracks, door
sills, cabinets, closets, around pipe openings and other
hiding places.

Spiders: Spray webs and corners where spiders gather.

Brown Dog Ticks: Spray or paint baseboards, door sills,
edges of carpet, behind pictures and draperies, and all
other possible hiding places. **DO NOT TREAT DOGS
WITH THIS PRODUCT.**

Wasps: Spray nests, particularly nest openings; around
windows and areas frequented by wasps.

Boxelder Bugs: Spray around or side of buildings where
insects crawl. Repeat application as needed.

WARNINGS: Use on limited areas only. Do not treat
entire walls or floor. Avoid spotting wallpaper and
fabrics. Do not contaminate food, water, dishes or
utensils. Allow treated dish and food shelves to dry;
cover with shelf paper before replacing articles. Do not
allow children, pets or animals near treated surfaces
until spray has dried. Do not use in dairy barns or
poultry houses.

This product is toxic to fish and wildlife. Keep out of
any body of water. Apply this product only as specified
on this label.