



E P A REG. NO. 602-176-AA



# PURINA® RABON\* INSECTICIDE

**50% WETTABLE POWDER**



**PURINA RABON INSECTICIDE**

is suitable for use in  
conventional hydraulic  
sprayers or low  
pressure equipment.

**RESIDUAL FLY CONTROL AND MAGGOT KILLER  
AND CONTROLS NORTHERN FOWL MITES, CHICKEN  
MITES AND LICE ON OR AROUND CHICKENS**

ACTIVE INGREDIENTS

2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate . . . . .	50.0%
INERT INGREDIENTS . . . . .	50.0%
	100.0%

\*Reg. T.M. Shell Chemical Company

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE OTHER CAUTIONS ON BACK PANEL**

**NET WEIGHT 4 LB.**

*Manufactured by*

**RALSTON PURINA COMPANY**  
General Offices • Checkerboard Square • St. Louis, Missouri 63188

6-26-72

602-176



**HOW TO USE**

**PURINA RABON INSECTICIDE 50% WETTABLE POWDER**

**MIXING DIRECTIONS FOR STRENGTH AND AMOUNT NEEDED**

Strength	1 gallon	3 gallons	25 gallons	100 gallons
½%	2½ Tbls.*	¼ lb.	2 lbs.	8 lbs.
1%	5 Tbls.	½ lb.	4 lbs.	16 lbs.

Note: Agitate mixture thoroughly before application. \*Tablespoonful

**FLY CONTROL**

**FARM BUILDING SPRAY**

Area	Strength	Application
Dairy, Poultry, Hog, Beef Cattle, Horse and Other Farm Buildings	1%	Apply thoroughly to point of runoff to walls, ceilings & where flies congregate and feed. Use 1 gallon for each 500-1000 square feet. Repeat as necessary. Do not apply this strength spray to birds or animals.

**MAGGOT CONTROL**

Area	Strength	Application
Caged Poultry, Manure Piles and Other Fly Breeding Areas	1%	Apply 1 gallon of mixture to every 100 square feet of droppings or surface area. Repeat as necessary.

**POULTRY LICE AND MITE CONTROL**

Area	Strength	Application
Caged Hens	½%	Apply directly 1 gallon to 100 birds using power sprayer at 100 psi. For individual bird treatment carefully apply 1 ounce per bird. Spray vent & fluff areas from below. Repeat when necessary but not more often than every 14 days.
Floor Birds	½%	Apply 1 to 2 gallons per 1000 sq. ft. evenly to litter surface. Spray walls, roosts, cracks, crevices and interiors thoroughly. Spray birds lightly; treat roosters carefully to avoid reinfestation of breeding flocks.
Litter	50% W.P.	Apply 1 lb. to 640 sq. ft. Treat evenly and thoroughly.
Dust Box	50% W.P.	Use 2.5 oz. (15 level tablespoonfuls) per 50 birds. Mix evenly throughout top layer of box content.
Roost Paint	1%	Use 1 pint to 100 feet. Apply thoroughly with brush or sprayer, especially to cracks.
Fowl Tick (Blue Bug)	1%	Use 1 gallon for 100 to 150 sq. ft. Apply thoroughly to walls, ceilings, floor cracks and crevices with power sprayer.



**PURINA RABON INSECTICIDE**  
is suitable for use in  
conventional hydraulic  
sprayers or low  
pressure equipment.



**PRECAUTIONS IN USING**

May be fatal if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

**QUALITY CONTROLLED BY PURINA RESEARCH**