

Reason for Issue: Warning statement
reworded. First aid statement added.

Print Date Released 3-1-73

Date Released _____

Front Panel
ORIGINAL

RAVAP* INSECTICIDE
for
RESIDUAL FLY CONTROL AND MAGGOT KILLER

E.P.A. Reg. No. 602-177-AA

(R) In Checkers)

contains

2 lb. RABON* Insecticide and 0.5 lb. VAPONA* Insecticide per gallon
Emulsifiable

ACTIVE INGREDIENTS

2-chloro-1-(2,4,5-trichlorophenyl)	
vinyl dimethyl phosphate	23.0%
2,2-dichlorovinyl dimethyl phosphate	5.3%**
Related Compounds	0.4%**
Xylene	48.3%
INERT INGREDIENTS	23.0%
	<hr/> 100.0%

*Reg. T.M. Shell Chemical Company
**Equivalent to 5.7% w. Vapona* Insecticide



OBSERVE LABEL DIRECTIONS

Net Contents: 5 Gallons

WARNING: KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS

SEE AFTERTHOUGHT AND OTHER PRECAUTIONS ON THE PANEL

REG. U.S. PAT. & TM. OFF.

(Note: Type size and
prominence will conform
with precautionary
labeling requirements).

Manufactured by
Ralston Purina Company
General Office
Checkerboard Square
St. Louis, Missouri 63108

E.P.A. Approved
3-16-73
Ready to print
Ready to market

Side Panel

How To Use Purina RAVAP Insecticide

Purina RAVAP Insecticide is suitable for use in conventional hydraulic sprayers or low pressure knapsack sprayers.

USE AND APPLICATION DIRECTIONS

RESIDUAL SURFACE SPRAY: Mix 1 gallon of RAVAP Insecticide in 25 gallons of water. Apply as an over-all application to ceilings and walls of dairy barns, livestock sheds, calf barns, hog barns and other farm buildings. Do not use RAVAP in poultry houses as a residual surface spray. Other areas where flies congregate, such as around garbage dumps and corrals, should be sprayed thoroughly. Repeat applications as fly populations warrant. Extreme infestations may necessitate increasing the dosage to 2 gallons RAVAP Insecticide per 25 gallons of water. Applications should be made to the point of run-off. One gallon of spray should treat 500 to 1000 square feet of surface area.

MAGGOT KILLER: Mix 1 gallon of RAVAP Insecticide with 25 gallons of water. Apply 1 gallon of solution for every 100 square feet of droppings under poultry on wire or in cages; or apply around manure or garbage. Do not apply to poultry.

DILUTION TABLE

RAVAP - 6 oz.	1 pint	1 gal.
WATER - 1 gal.	3 gals.	25 gals.

DANGER: Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly, a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Apply this product only as specified on this label. Do not contaminate feed or food products. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Rinse equipment and containers; dispose of wastes by burying in non-crop lands away from water supplies. Do not re-use empty containers. Destroy by burying with wastes or by burning. Stay away from smoke or fumes.