PURINA LAWN WEED KILLER

CONTROLS: Dandelion, Chickweed, Plantain, Henbit, Knotweed, Purslane, Oxalis, Wild Onion, Carpetweed, Curly Dock, Clover, Sheep Sorrel, Thistles and certain other broad-leaf weeds.

DIRECTIONS: Spray when weeds are actively growing. Apply when air is calm to avoid drift onto desirable plants. Spray entire lawn or spot treat as needed. During dry periods, water lawn thoroughly 1 or 2 days before using Lawn Weed Killer. After using, do not sprinkle for 24 hours.

Repeat applications in 2 to 3 weeks on hard-to-kill weeds. Treat Chickweed, Henbit and Wild Onion in Fall and Spring if needed.

HOW TO MIX: (Make new dilution for each use; do not store diluted spray.) **For Spot Spraying**—mix 1 tablespoonful to 1 gallon of water. Spray each weed to point of runoff.

For Lawn Spraying—Determine size of lawn area.

Use ½ ounce (1 tablespoonful) in 1 gallon of water for each 200 sq. ft. of hardy lawn grass.

Use ¼ ounce (½ tablespoonful) in 1 gallon of water for each 200 sq. ft. of Bent grass, St. Augustine, Centipede, Carpet or Red Fescue. Temporary injury may occur on these grasses; make two applications 10 days apart.

NOTE: Do not treat newly seeded lawns until grass is well established. Do not reseed for 4 weeks after treatment. Do not apply to Dichondra, Lippia or where desirable clover is present.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Wash hands after use. Keep out of the reach of children.

Do not use for controlling weeds where vapors may injure susceptible plants. Coarse sprays are less likely to drift. Do not contaminate domestic or irrigation water.

Do not store near feed, foodstuffs, fertilizers, seeds, insecticides or fungicides. When container is empty wash thoroughly and destroy. Do not use same spray equipment for other purposes.

QUALITY CONTROLLED BY PURINA RESEARCH

706A   G-5253   Printed in U.S.A.

U.S.D.A. Reg. No. 602-180

PURINA®
LAWN WEED KILLER

FOR LAWN USE ONLY

CONTAINS 2, 4-D AND SILVEX

TAKE TIME

OBSERVE LABEL DIRECTIONS

Active Ingredients
*Isooctyl Ester of 2,4-Dichlorophenoxyacetic acid 22.90%
**Isooctyl Ester of Silvex [2-(2,4,5-Trichlorophenoxy) propionic acid] . 10.75%
Inert Ingredients 66.35%
                                                                    100.00%

*Equivalent to 15.2% 2,4-Dichlorophenoxyacetic acid
**Equivalent to 7.6% Silvex [2-(2,4,5-Trichlorophenoxv) propionic acid].

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS: 16 FL. OZ. (1 PINT)

Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63199