

EPA Reg No 602 183



# FRUIT TREE SPRAY

Controls Insects and Plant Diseases of  
Fruits and Berries



#### Active Ingredients

*Malathion	8.33%
**Methoxychlor, Technical	16.67%
***Captan	16.67%
Inert ingredients	58.33%
	100.00%

\* Malathion - 10% granulated, 20% liquid, 40% dust.

\*\* Methoxychlor - 44% technical, 20% dust.

\*\*\* Captan - 70% dust, 30% granulated, 10% liquid.

\*\*\*\* Proprietary blend of various inert ingredients.

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE OTHER WARNINGS ON BACK PANEL**

NET WT. 48 OZ. (3 LB.)

## DIRECTIONS FOR USE

Mix 4 level tablespoonfuls of Fruit Tree Spray per gallon of water (1 lb. to 16<sup>2</sup> gallons). Pre mix required amount of Fruit Tree Spray with a little water to make a thin slurry before adding all the water. Keep mixture agitated while spraying. Always use freshly mixed solution.

Spray upper and lower leaf surfaces and fruit thoroughly. Apply 2 to 3 gallons of spray per medium sized tree (6-8 feet high); larger trees may require 3 to 6 gallons or more.

**Apples, Pears** - Aphid, apple maggot, cankerworm, codling moth, plum curculio, eye-spotted bud moth, European red mite, Japanese beetle, leafhopper, leaf miner, leaf roller, scale (Forbes, Putnam, and San Jose species), two spotted mite. **Diseases** - Scab, bitter rot, sooty blotch, fly speck, cedar apple rust.

Spray trees when flower buds first begin to separate. Repeat at 7 day intervals when buds show pink. Do not spray when blossoms are open. After petal fall, repeat every 14 days. Do not spray within 7 days of harvest.

**Peaches, Plums** - Aphid, plum curculio, Japanese beetle, Oriental fruit moth, peach twig borer, rose chafer, scale (cottony peach, European fruit, Forbes, San Jose and terrapin species), tarnished plant bug. **Diseases** - Scab, brown rot, blossom blight.

Spray trees just before blossoms open. Do not spray when blossoms are open. Respray when most petals have dropped and at 10 day intervals thereafter. Do not spray peaches within 21 days or plums within 7 days of harvest.

**Cherries, Apricots** - Aphid, plum curculio, cherry fruitworm, cherry fruit fly, Japanese beetle, red spider mite, rose chafer, thrip, Forbes scale, San Jose scale, slug. **Diseases** - Leaf spot, brown rot, blossom blight, scab.

Spray trees when most of the petals have fallen. Respray at 10 day intervals. Do not spray cherries within 7 days of apricots within 21 days of harvest.

**Raspberries, Blackberries** - Aphid, leafhopper, Japanese beetle, rose chafer, rose scale. **Diseases** - Anthracnose, spur blight.

Spray bushes when insects are first seen; repeat as needed for control. For diseases apply when blossoms are in bud (young canes 8 to 10 inches long). Respray 1 week later and in fall when old canes are removed. Do not apply within 14 days of harvest.

**Grapes** - Grapes - aphid, leafhopper, grape berry moth, Japanese beetle, rose borer, two spotted spider mite. **Diseases** - Black rot, downy mildew.

Spray vines when insects first appear. Respray at 10 day intervals as needed. For diseases apply just before bloom, repeat immediately after bloom (1 to 3 more sprays may be needed). Do not spray within 14 days of harvest.

**Strawberries** - Aphids, slugs, beetles, mites, thrip, Japanese beetle, spider mites, spittlebug, strawberry leafroller, strawberry leafroller, strawberry weevils, white fly, weevils, thrip. **Diseases** - Botrytis fruit rot.

Spray plants when insects first appear. Respray at 10 day intervals as needed. Do not spray when flowers are open. Do not spray to form regular spray. Do not spray within 14 days of harvest.


**CAUTION:** Harmful to bees. Do not spray fruit trees during bloom. Avoid drift to blossoming trees. Wash hands thoroughly after use. Avoid contact with eyes and skin.

**ANTIDOTE:** If swallowed, drink a glass of warm water. Do not induce vomiting. If skin contact occurs, wash with soap and water. For eye contact, flush eyes with water. Get medical attention.

This product contains malathion. Keep out of reach of children and animals. Do not contaminate water.

KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS

**Controls Insects and Plant Diseases of  
Fruits and Berries**



**Active Ingredients:**

*Malathion.....	8.33%
**Methoxychlor, Technical.....	16.67%
***Captan.....	16.67%
Inert Ingredients.....	58.33%
	100.00%

\*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

\*\*Equivalent to 14.67% 2,2-Bis (p-methoxyphenyl),1,1,1-trichloroethane and 2.00% of other isomers and related compounds.

\*\*\*N-trichloromethylthio-4 cyclohexene-1,2-dicarboximide

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE OTHER WARNINGS ON BACK PANEL**

**NET WT. 48 OZ. (3 LB.)**

Manufactured by