

602-185

ACCT

Jul 1

CHLORINE SANITIZER 0-10

WARNING: Handle with care. When measuring PURINA CHLORINE SANITIZER 0-10, use only a china, plastic, or enamel cup or spoon that is completely dry, clean, and free from any foreign matter, liquid, or solid. PURINA CHLORINE SANITIZER 0-10 is a strong oxidizing agent. Contact with heat, acids, combustible organic materials, such as vinegar, paint products, beverages, tobacco, cleaners, soap products, pine oil, kerosene, paper, rags, etc., may cause fire. In case of fire, douse liberally with water. Never mix PURINA CHLORINE SANITIZER 0-10 with anything except water. Keep in a cool dry place in original container. Always replace lid.

DO NOT REUSE CONTAINER, DESTROY WHEN EMPTY

DIRECTIONS:

1. To sanitize table tops and counters in food preparation areas, first thoroughly clean area to be sanitized. Then prepare a sanitizing solution by dissolving 1 ounce of PURINA CHLORINE SANITIZER 0-10 in 4 gallons of water (200 ppm available chlorine). Apply sanitizing solution with a clean cloth so as to wet surface thoroughly. Rinse with potable water.

2. To sanitize equipment in food processing plants and dairies, first thoroughly clean area to be sanitized. Then prepare a sanitizing solution by dissolving 1 ounce of PURINA CHLORINE SANITIZER 0-10 in 4 gallons of water (200 ppm available chlorine). Apply sanitizing solution with a clean cloth or through mechanical equipment so as to wet surface thoroughly. Rinse with potable water.

POISON

WARNING: Harmful if swallowed. May cause chemical burns. Do not allow contact with skin, mucous membranes, or clothing.

ANTIDOTE:

External: Flood with water for 15 minutes.


Internal: Drink large quantities of water or milk. Follow with milk of magnesia, vegetable oil, or beaten eggs. Call a physician immediately.

Eyes: Wash eyes with water for 15 minutes. Get prompt medical attention.

KEEP IN A COOL DRY PLACE—REPLACE LID TIGHTLY
FOR FURTHER ASSISTANCE CONTACT YOUR PURINA REPRESENTATIVE
QUALITY ASSURED BY PURINA RESEARCH

Purina

CHLORINE SANITIZER 0-10



POISON

U.S.D.A. Reg. No. 602-185

Active Ingredient:	
Sodium dichloro-s-triazinetrione.....	17.50%
Inert Ingredients.....	82.50%
Available Chlorine	10%

**CHEMICAL
SPECIALTIES**

WARNING: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER REQUIRED WARNING
STATEMENTS ON BACK PANEL

Keep Lid Tightly Closed When Not In Use
Mix Only With Water
May Cause Fire If Contaminated

NET WT. 5 LB.

S-1660

→ A chlorine sanitizer, deodorant,
and destainer for Food Processing
Plants, Dairies, Restaurants, and
Institutions

**READILY SOLUBLE
EASY TO USE
SANITIZES
DESTAINS
DEODORIZES**



Manufactured by
RALSTON PURINA COMPANY
General Office • Heckert Building • St. Louis, Mo. 63188

602-185

ACCEPTED

Jul 11 1954

3

PURINA CHLORINE SANITIZER 0-10

WARNING: Handle with care. When measuring PURINA CHLORINE SANITIZER 0-10, use only a china, plastic, or enamel cup or spoon that is completely dry, clean, and free from any foreign matter, liquid, or solid. PURINA CHLORINE SANITIZER 0-10 is a strong oxidizing agent. Contact with heat, acids, combustible organic materials, such as vinegar, paint products, beverages, tobacco, cleaners, soap products, pine oil, kerosene, paper, rags, etc., may cause fire. In case of fire, douse liberally with water. Never mix PURINA CHLORINE SANITIZER 0-10 with anything except water. Keep in a cool dry place in original container. Always replace lid.

DO NOT REUSE CONTAINER, DESTROY WHEN EMPTY

DIRECTIONS:

1. To sanitize table tops and counters in food preparation areas, first thoroughly clean area to be sanitized. Then prepare a sanitizing solution by dissolving 1 ounce of PURINA CHLORINE SANITIZER 0-10 in 4 gallons of water (200 ppm available chlorine). Apply sanitizing solution with a clean cloth so as to wet surface thoroughly. Rinse with potable water.

2. To sanitize equipment in food processing plants and dairies, first thoroughly clean area to be sanitized. Then prepare a sanitizing solution by dissolving 1 ounce of PURINA CHLORINE SANITIZER 0-10 in 4 gallons of water (200 ppm available chlorine). Apply sanitizing solution with a clean cloth or through mechanical equipment so as to wet surface thoroughly. Rinse with potable water.

POISON

WARNING: Harmful if swallowed. May cause chemical burns. Do not allow contact with skin, mucous membranes, or clothing.

ANTIDOTE:

External: Flood with water for 15 minutes.
Internal: Drink large quantities of water or milk. Follow with milk of magnesia, vegetable oil, or beaten eggs. Call a physician immediately.
Eyes: Wash eyes with water for 15 minutes. Get prompt medical attention.

KEEP IN A COOL DRY PLACE—REPLACE LID TIGHTLY
FOR FURTHER ASSISTANCE CONTACT YOUR PURINA REPRESENTATIVE
QUALITY ASSURED BY PURINA RESEARCH

7131 Printed in U.S.A.

PURINA CHLORINE SANITIZER



POISON

U.S. Active Ingre Sodium Inert Ingre Available Cr

WARNING: KEEP LID TIGHTLY CLOSED. SEE ANTIDOTE AND STATEMENT.

Keep Lid Tight
Mix
May Cause

NET

DOCUMENT AVAILABLE

15

3

ment in food processing plants
highly clean area to be sanitized.
rizing solution by dissolving 1
LORINE SANITIZER 0-10 in 4
ppm available chlorine). Apply
with a clean cloth or through
so as to wet surface thoroughly.
eter.

allowed. May cause chemical
contact with skin, mucous mem-

water for 15 minutes.
quantities of water or milk. Follow
p, vegetable oil, or beaten eggs.
mediately.
water for 15 minutes. Get prompt

PLACE-REPLACE LID TIGHTLY
CONTACT YOUR PURINA REPRESENTATIVE
BY PURINA RESEARCH

Printed in U.S.A.

15

Purina

CHLORINE SANITIZER 0-10



U.S.D.A. Reg. No. 602-185



POISON

Active ingredient:
Sodium dichloro-s-triazinetrione..... 17.50%
Inert ingredients..... 82.50%
Available Chlorine 10%

**CHEMICAL
SPECIALTIES®**

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER REQUIRED WARNING
STATEMENTS ON BACK PANEL**

Keep Lid Tightly Closed When Not In Use
Mix Only With Water
May Cause Fire If Contaminated

NET WT. 5 LB.

BEST DOCUMENT AVAILABLE

4

S-1660

→ A chlorine sanitizer, deodorant,
and destainer for Food Processing
Plants, Dairies, Restaurants, and
Institutions / PREMISES

**READILY SOLUBLE
EASY TO USE
SANITIZES
DESTAINS
DEODORIZES**

TAKE TIME



Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63103

16