

HOW TO USE PURINA DAIRY SPRAY SPECIAL

Purina Dairy Spray Special is a ready-to-use oil base spray for use in hand sprayers, misting equipment, and back or face rubbers. Contains Ciodrin* and Vapona* Insecticides for effective knock-down, kill and residual protection against flies and lice on dairy or beef cattle.

For Control of House Flies, Stable Flies, Horn Flies, and Face Flies: Apply Dairy Spray Special as a mist to the hair coat by using a hand atomizer-type sprayer or mechanical misting equipment. Spray once daily using 1 to 2 ounces over the entire animal. Spray calves more sparingly. Do not wet the skin.

For stable flies, spray legs and flanks; for face flies spray face and head areas. Repeat once daily as necessary.

Back Rubbers and Face Rubbers: For the control of horn flies and face flies on lactating dairy and beef animals:

Use 1 gallon of Dairy Spray Special for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

For Control of Lice: Make 1 to 2 applications of 3-3½ ounces of Dairy Spray Special at 14 day intervals. Apply as a fine mist spray to thoroughly cover all parts of the animal. If applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

CAUTION: Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Wash thoroughly with soap and water after using, and before eating or smoking. Harmful or fatal if swallowed. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.



Keep out of reach of children. Do not mist or fog into air of closed buildings. Do not wet the skin of animals. Do not spray or store near an open flame. Do not contaminate feed, water or foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

QUALITY CONTROLLED BY PURINA RESEARCH

712J

Printed in U.S.A.




4-17-77
602-186



Net Contents: 1 Gallon D-2565

U.S.D.A. Reg. No. 602-186

PURINA[®] DAIRY SPRAY SPECIAL



CAUTION: KEEP OUT OF REACH OF CHILDREN
See Other Cautions on Back Panel

READY-TO-USE OIL BASE FLY SPRAY

Kills House Flies, Stable Flies, Face Flies, Horn Flies, Lice on Dairy or Beef Cattle.

Active Ingredients:

Dimethyl Phosphate of Alpha methylbenzyl-3 Hydroxy-cis Crotonate	1.00%
2,2-Dichlorovinyl Dimethyl Phosphate**.....	0.23%
Related Compounds**.....	0.02%
Petroleum Oils.....	98.50%
Inert Ingredients.....	0.25%
	100.00%

**Equivalent to 0.25% Vapona Insecticide

Manufactured by

RALSTON PURINA COMPANY

General Offices • St. Louis, Missouri 63188

**NEW
FORMULA**



1-12-77
6-11-77

CIAL

ations of 3-3 1/2 ounces of
ils. Apply as a fine mist
the animal. If applied by
imum spraying time per

contact with skin. In case
plenty of water, for eyes
with soap and water after
mful or fatal if swallowe
ly and get medical atten
so antidotal and may be
le.

st or fog into air of closed
ls. Do not spray or store
eed, water or foodstuffs.

Birds feeding on treated
ly of water. Do not apply
taminate water by clear
apply this product onl