

1 of 2

ACCEPTED
MAY 25 1977
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, and registered under the pesticide EPA Reg. No. 602-186




PURINA® DAIRY SPRAY SPECIAL

E P A Reg. No. 602-186-AA



READY-TO-USE OIL BASE FLY SPRAY

Kills House Flies, Stable Flies, Face Flies, Horn Flies, Lice on Dairy or Beef Cattle.



Active Ingredients:

Dimethyl Phosphate of Alpha-methylbenzyl 3-Hydroxy-cis-Crotonate.....	1.00%
2,2-Dichlorovinyl Dimethyl Phosphate**.....	0.23%
Related Compounds**.....	0.02%
Petroleum Oils.....	98.50%
Inert Ingredients.....	0.25%
	100.00%

**Equivalent to 0.25% Vapona Insecticide

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS BELOW

BEST AVAILABLE COPY

HOW TO USE PURINA DAIRY SPRAY SPECIAL

Purina Dairy Spray Special is a ready-to-use oil base spray for **For Control of Lice:** Make 1 to 2 applications of 3-3½ ounces of

PM 15
602-186

Active Ingredients:		
Dimethyl Phosphorothioate	of Alpha-methylbenzyl 3-Hydroxy-cis-Crotonate	1.00%
2,2-Dichlorovinyl	methyl Phosphate**	0.25%
Related Compounds		0.05%
Petroleum Oils		98.50%
Inert Ingredients		0.25%
		100.00%

**Equivalent to 0.25% Vapona Insecticide

BEST AVAILABLE COPY

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS BELOW**

HOW TO USE PURINA DAIRY SPRAY SPECIAL

Purina Dairy Spray Special is a ready-to-use oil base spray for use in hand sprayers, misting equipment, and back or face rubbers. Contains Ciodrin* and Vapona* Insecticides for effective knock-down, kill and residual protection against flies and lice on dairy or beef cattle. Do not apply on horses.

For Control of House Flies, Stable Flies, Horn Flies, and Face Flies: Apply Dairy Spray Special as a mist to the hair coat by using a hand atomizer-type sprayer or mechanical misting equipment. Spray once daily using 1 to 2 ounces over the entire animal. Do not wet the skin due to possible skin irritation. Do not spray calves under 1 month of age. Do not apply regularly to calves under 6 months of age.

For stable flies, spray legs and flanks; for face flies spray face and head areas. Repeat once daily as necessary.

Back Rubbers and Face Rubbers: For the control of horn flies and face flies on lactating dairy and beef animals:

Use 1 gallon of Dairy Spray Special for each 20 linear feet of back-up back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

For Control of Lice: Make 1 to 2 applications of 3-3½ ounces of Dairy Spray Special at 14 day intervals. Apply as a fine mist spray to thoroughly cover all parts of the animal. If applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

CAUTION: Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Wash thoroughly with soap and water after using, and before eating or smoking. Harmful or fatal if swallowed. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Keep out of reach of children. Do not mist or fog into air of closed buildings. Do not wet the skin of animals. Do not spray or store near an open flame. Do not contaminate feed, water or foodstuffs. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not re-use empty container. Destroy container by perforating and bury or discard in a safe place.

EPA Est. No. 602-MO-1

QUALITY CONTROLLED BY PURINA RESEARCH

Manufactured by

RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Missouri 63188

NET 39 GALS.

D-2568

Printed in U.S.A.

BEST AVAILABLE COPY