


PM 15  
602-186


1 of 4



# PURINA DAIRY SPECIAL

READY-TO-USE OIL

Kills House Flies, Stable Flies, Face Flies,



**Active Ingredients:**

Dimethyl Phosphate of Alpha-methylbenzyl  
2,2-Dichlorovinyl Dimethyl Phosphate\*\*  
Related Compounds\*\*  
Petroleum Oils

**Inert Ingredients:**

\*\*Equivalent to 0.25% Vapona Insecticide

**CAUTION: KEEP OUT  
SEE OTHER SIDE**

## HOW TO USE PURINA

Purina Dairy Spray Special is a ready-to-use oil base spray for use in hand sprayers, misting equipment and back or face rubbers. Contains Dieldrin\* and Vapona\* Insecticides for effective knock-down, kill and residual protection against flies and lice on dairy or beef cattle. Do not apply on horses.

**For Control of House Flies, Stable Flies, Horn Flies, and Face Flies:** Apply Dairy Spray Special as a mist to the hair coat by using a hand atomizer-type sprayer or mechanical misting equipment. Spray once daily using 1 to 2 ounces over the entire animal. Do not wet the skin due to possible skin irritation. Do not spray calves under 1 month of age. Do not apply regularly to calves under 6 months of age.

For stable flies, spray legs and flanks; for face flies spray face and head areas. Repeat once daily as necessary.

**Back Rubbers and Face Rubbers:** For the control of horn flies and face flies on lactating dairy and beef animals:

Use 1 gallon of Dairy Spray Special for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

**For Control of Lice:** Make 1 to 2 applications of 3-3½ ounces of Dairy Spray Special at 14 day intervals. Apply as a fine mist spray to thoroughly cover all parts of the animal. If applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

**CAUTION:** Avoid inhalation of mist and contact with skin. In case of contact with skin, or eyes, flush with plenty of water, for eyes get medical attention. Wash thoroughly with soap and water after

7750

**QUALITY CONTROLLED**

Manuf  
**RALSTON PURINA**  
General Offices • Checkerboard S

# NET 30 GAL.

2 of 4

# URINA<sup>®</sup> FLY SPRAY SPECIAL

EPA Reg. No. 602-186




10-USE OIL BASE FLY SPRAY

Flies, Face Flies, Horn Flies, Lice on Dairy or Beef Cattle.



2-Methylbenzyl 3-Hydroxy-cis-Crotonate.....	1.00%
Phosphate**.....	0.23%
.....	0.02%
.....	98.50%
.....	0.25%
.....	100.00%



KEEP OUT OF REACH OF CHILDREN  
OTHER CAUTIONS BELOW

## URINA DAIRY SPRAY SPECIAL

Use for use  
rubbers.  
knock-  
dairy or

Flies:  
hand  
Spray  
wet the  
under 1  
of age.  
face and

flies and

Burlap  
rvoir as  
control.  
of Dairy  
spray to  
ging or  
al is 20

In case  
for eyes  
ter after

using, and before eating or smoking. Harmful or fatal if swallowed. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Keep out of reach of children. Do not mist or fog into air of closed buildings. Do not wet the skin of animals. Do not spray or store near an open flame. Do not contaminate feed, water or foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or Local disposal authorities.

### CONTAINER DISPOSAL:

- (a) Reseal container and offer for reconditioning or
- (b) Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place.


Consult Federal, State or Local disposal authorities for approved alternative procedures.

ROLLED BY PURINA RESEARCH

Manufactured by

PURINA COMPANY

Merck Square • St. Louis, Missouri 63188



EPA Est. No. 602-MO-1

D-2568

Printed in U.S.A.


**BEST AVAILABLE COPY**

BEST AVAILABLE COPY



# PURINA® DAIRY SPRAY SPECIAL

READY-TO-USE OIL-BASE FLY SPRAY



**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE OTHER CAUTIONS ON BACK PANEL**

Kills House Flies, Stable Flies, Face Flies, Horn Flies, Lice on Dairy or Beef Cattle.

Active Ingredients:	
4-Dimethyl Phosphate of Alpha-methylbenzyl	1.00%
3-hydroxy-cis-Crotonate	0.23%
2,2-Dichlorovinyl Dimethyl Phosphate*	0.02%
Related Compounds**	98.50%
Petroleum Oils	0.25%
Inert Ingredients	100.00%
**Equivalent to 0.25% Vapona Insecticide	

## NET CONTENTS: 1 GALLON

Manufactured By  
**RALSTON PURINA COMPANY**  
General Offices • Checkerboard Co. • St. Louis, Mo. 631

BEST AVAILABLE COPY

## HOW TO USE PURINA DAIRY SPRAY SPECIAL

Purina Dairy Spray Special is a ready-to-use oil base spray for use in hand sprayers, misting equipment and back or face rubbers. Contains Ciodrin\* and Vapona\* Insecticides for effective knock-down, kill and residual protection against flies and lice on dairy or beef cattle. Do not apply on horses.

**For Control of House Flies, Stable Flies, Horn Flies, and Face Flies:** Apply Dairy Spray Special as a mist to the hair coat by using a hand atomizer-type sprayer or mechanical misting equipment. Spray once daily using 1 to 2 ounces over the entire animal. Do not wet the skin due to possible skin irritation. Do not spray calves under 1 month of age. Do not apply regularly to calves under 6 months of age.

For stable flies, spray legs and flanks; for face flies spray face and head areas. Repeat once daily as necessary.

**Back Rubbers and Face Rubbers:** For the control of horn flies and face flies on lactating dairy and beef animals:

Use 1 gallon of Dairy Spray Special for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

**For Control of Lice:** Make 1 to 2 applications of 3-3 1/2 ounces of Dairy Spray Special at 14 day intervals. Apply as a fine mist spray to thoroughly cover all parts of the animals. If applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

**CAUTION:** Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get

medical attention. Wash thoroughly with soap and water after using, and before eating or smoking. Harmful or fatal if swallowed. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Keep out of reach of children. Do not mist or fog into air of closed buildings. Do not wet the skin of animals. Do not spray or store near an open flame. Do not contaminate feed, water or foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

### STORAGE AND DISPOSAL

#### PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

#### PESTICIDE DISPOSAL

Pesticide, spray, mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

#### CONTAINER DISPOSAL

Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill or bury in a safe place.

Consult Federal, State or Local disposal authorities for approved alternative procedures.

QUALITY CONTROLLED BY PURINA RESEARCH