U.S.D.A. Reg. No. 602-187

NEW FORMULA

TAKE TIME

OBSERVE LABEL DIRECTIONS

PURINA®

Kills... Horn Flies, House Flies, Mosquitoes, Gnats, and Lice.

Active Ingredients:

Pyrethrins..... Piperonyl butoxide, technical*..... 0.48% Petroleum Oils..... 99.46%

*Consists of 0.384% of (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.096% of Related Compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN See Other Warnings on Back Panel.

NET CONTENTS: 1 GALLON

Manufactured by RALSTON PURINA COMPANY

General Offices • Checkerboard Square • St. Louis, Missouri 63188











HOW TO USE PURINA DAIRY SPRAY

Purina Dairy Spray is a quick knockdown contact spray for killing and repelling horn flies, house flies, mosquitoes and gnats and also kills lice. For fly control, apply daily; after using three days the residual repellent and killing action may be sufficient so that applications may be needed only every other day.

Dairy Cattle: Fly Control—Apply Purina Dairy Spray as a mist to the hair coat. Hold sprayer about 18 inches from the cow; apply mist to entire body from mid-neck region to tail, including the underline and the legs. The amount of spray used should not exceed 1 to 2 ounces per adult animal. Spray calves more sparingly. Do not wet the skin. Wash udder before milking.

Lice Control—Spray the animals with 2 ounces of Purina Dairy Spray. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

Dairy Barn and Milk Room. When used as a space contact spray for flies and other insects, remove animals from barn and close doors and windows tightly. Spray Purina Dairy Spray towards ceiling until a light fog hangs in the air. Repeat as necessary.

Horse, Beef Cattle, Hogs, and Other Farm Animals—Apply as a mist to the hair coat of farm animals for temporary repellent and killing action. Aids in controlling horn flies, house flies, mosquitoes and gnats. Do not wet the skin.

CAUTION: Harmful or fatal if swallowed. If swallowed do not induce vomiting. Call physician immediately. Avoid breathing of spray mist.

Do not use, pour, spill or store near an open flame. Do not wet the skin of animals. Avoid contamination of feed, foodstuffs, milk and milking utensils. Remove animals from barn when fogging or misting.

Do not re-use empty container. Wash thoroughly with water and detergent. Discard in a safe place.

714G

D-2495

Printed in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH





